

Research Article

# Biodiversity and the Community Structure of Chromista Cavalier-Smith, 1981 in Nyong and Kienke River Mouths (South-Cameroon)

**Christelle Chimène Mokam<sup>1,2</sup>** , **Andrea Sarah Kenne Toukem<sup>1</sup>** ,  
**Christian Dongmo Teufack<sup>3</sup>** , **Fabien Trésor Amougou Dzou<sup>3</sup>** ,  
**Sedrick Junior Tsekane<sup>1</sup>** , **Mohammadou Moukhtar<sup>1</sup>** ,  
**Auguste Pharaon Mbianda<sup>4</sup>** , **Martin Kenne<sup>1,\*</sup>** 

<sup>1</sup>Department of Biology and Physiology of Animal Organisms, Faculty of Science, University of Douala, Douala, Cameroon; Department of Biology, Ecology and Evolution, Liège (Sart-Tilman), Belgium

<sup>2</sup>Department of Biology, Ecology and Evolution, Liège (Sart-Tilman), Belgium

<sup>3</sup>Laboratory of the Specialized Center for Research on Marine Ecosystems, Kribi, Cameroon

<sup>4</sup>Department of Biology of Vegetal Organisms, Faculty of Science, University of Douala, Douala, Cameroon

## Abstract

A survey was undertaken from March to June 2014 on the biodiversity and the community structure of Chromista Cavalier-Smith, 1981 in Nyong and Kienke River mouths (South-Cameroon). In each river, raw waters were collected from upstream to downstream at four sites. Cells were counted using the Malassez cells procedure and species were identified. A total of  $10427.1 \times 10^5$  cells corresponded to three phyla, eight classes, 23 orders, 32 genera and 40 species (24 freshwater species (60.0% of total species richness and total collection respectively), three marine species (7.5% and 2.4% of the total species richness; and total collection respectively), and one brackish water specialist in Kienke (2.5% and 5.1%), 13 tolerant species (32.5% and 32.6%)). The trophic diatom index revealed undisturbed conditions with no or little alteration of human origin and a low organic pollution (oligotrophic or mesotrophic state) (Nyong: TDI=52.7; Kienke: TDI=69.7; pooled assemblage: TDI=65.0). A low species richness was detected (richness ratio in Nyong:  $d=0.008$ ; Kienke:  $d=0.003$ ; pooled rivers:  $d=0.004$ ), a high species diversity (Shannon index close to maximum) (Nyong:  $H'=2.742$  and  $H'_{\max}=2.996$ ; Kienke:  $H'=2.685$  and  $H'_{\max}=2.996$ ; pooled rivers:  $H'=3.245$  and  $H'_{\max}=3.689$ ), a very low dominance by a few species (Berger-Parker index close to 0) (Nyong:  $I_{BP}=0.156$ ; Kienke:  $I_{BP}=0.175$ ; pooled rivers:  $I_{BP}=0.134$ ), and Hill's ratio were close to 1 (Nyong: Hill=0.819; Kienke: Hill=0.803; pooled rivers: Hill=0.722). The community was highly even with a high value of the Pielou's evenness close to 1 (Nyong:  $J=0.915$ ; Kienke:  $J=0.896$ ; pooled rivers:  $J=0.880$ ). Two useful species and one harmful species to fish were rare in Kienke. Species exhibited in Kienke and pooled data in rainy season, a positive global net association while it was negative in Nyong. Assemblage fitted Preston's model in Nyong with a high environmental constant in the dry season ( $m'=1.469$ ), low constant in the rainy season ( $m'=0.947$ ) and the pooled seasons ( $m'=0.853$ ). In Kienke constants were low (dry season:  $m'=0.574$ ; rainy season:  $m'=0.566$ ; pooled seasons:  $m'=0.581$ ) suggesting a evolved community in less disturbed environments where the majority of species showed moderate abundances. In the dry season, the pooled assemblage functionned on the basis of maintaining a complex information network (close to ecological balance) developed at spatio-temporal scales (ZM model) and it presented a low force of regeneration (fractal dimension of the distribution of individuals among species ( $1/\gamma=0.925<1$ ). The evolved oligotrophic state (close to natural

\*Corresponding author: medoum68@yahoo.fr (Martin Kenne), medoum1968@gmail.com (Martin Kenne)

**Received:** 1 March 2024; **Accepted:** 18 March 2024; **Published:** 2 April 2024



Copyright: © The Author(s), 2023. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

balance) of the chromists' community should be preserved and protected and the studied rivers classified as reference.

## Keywords

Freshwater Species, Microalgae, Species Composition, Useful Species, Assemblage Functioning, Water Quality

## 1. Introduction

According to the AlgaeBase, 50,589 species of living algae and 10,556 fossil species are documented, referred to four kingdoms (Eubacteria Woese & Fox, 1977; Chromista Cavalier-Smith, 1981; Plantae Haeckel, 1866; and Protozoa Goldfuss, 1818), 14 phyla, and 63 classes [1]. Algae are the third most speciose grouping of plant-like after the flowering plants ( $\approx 382,000$  species) and fungi ( $\approx 170,000$  species, including lichens) [1]. The most species-rich phylum is the golden and brown algae Heterokontophyta Moestrup, 1992 with 18 classes and 21,052 living species dominated by the diatoms class Bacillariophyceae Dangeard, 1933 with 18,673 species (16,427 living species and 2,239 fossil) [1]. The next most species-rich phyla are the red algae Rhodophyta Wettst., 1901 (7,276 living species), the oxygenic photosynthetic green algae (6,851 living species), the blue-green algae (Cyanobacteria Stanier, 1973 or Cyanobacteriota Oren *et al.* 2022: 5,723 living species), the charophytes (4,950 living species, including Charophyceae with 511 living species, and Zygnematophyceae with 4,335 living species), Dinoflagellata (2956 living species, including Dinophyceae, 2,828 species), and haptophytes (Haptophyta: 1,722 species with 517 living species) [1]. Chromista is distinguished from Plantae and other phytoplankton because of its more complex chloroplast-associated membrane topology and rigid tubular multipartite ciliary's hairs [2]. Chromista (colored algae) possess a brown plastid resulting from a primary or secondary endosymbiosis with red algae [3]. They represent one of the five eukaryotic kingdoms in the seven-kingdom classification of life [3-9]. Nowadays Chromista comprises six divisions [golden or golden-brown algae Chrysophyta Pascher (1914); red algae Cryptophycophyta; Haptophyta Caval.-Sm., 1986; brown algae Phaeophyta Kjellman, 1891; dinoflagellates Pyrrophytaphyta; and yellow-green algae Xanthophyta Al-lorge, 1930, emend. Fritsch, 1935]. It comprises seven classes, 53 orders, 151 families, 487 genera, 2,029 species, one subspecies and 138 varieties [10]. Chrysophyta includes one class (Chrysophyceae Pascher, 1914), 10 orders, 23 families, 74 genera, 288 species, and 36 varieties [10]. Cryptophycophyta includes one class, six families, 14 genera, 83 species, and four varieties [10]. Haptophyta includes one class, three orders, 10 families, 45 genera, and 90 species [10]. Phaeophyta includes one class, 12 orders, 35 families, 156 genera, 498 species, and 51 varieties [10]. Pyrrophytaphyta includes two classes, 20 orders 57 families, 145 genera, 913 species,

one subspecies, and 39 varieties [10]. Xanthophyta includes about 600 species and many of the 100 known genera contain only a few species [6]. It includes one class (Xanthophyceae), seven orders, 20 families 53 genera 157 species, and eight varieties, [10]. Based on the number of families, the descending ranking of the well-represented classes is as follows: the diatoms Bacillariophyceae (92 families; 3,399 species), Dinophyceae (55 families; 911 species), Phaeophyceae (35 families; 498 species), Chrysophyceae (23 families; 288 species), Xanthophyceae (20 families; 157 species), Prymnesiophyceae (10 families; 90 species), Cryptophyceae (six families; 83 species), Chloromonadophyceae (two families; two species), and Cyanophyceae (more than 150 genera; 7,500 species). But these numbers are clearly below reality because the group has remained unknown to scientists for a long time and several forms are undetermined [7]. Chromists (colored algae) include the majority of marine species, brackish water specialists, freshwater species and terrestrial species as well as heterotrophic protists whether marine, brackish or freshwater. For these reasons, the Chromista kingdom is very important for the ocean and freshwater ecology [11]. As photosynthetic species they are essential primary producers of the aquatic food webs for fish and other macro invertebrates as well as microvertebrates, and they also have an economic importance as oxygen producers [12], or as bioindicators of the water quality (due to the preference of each species for aquatic environments with very specific physicochemical characteristics) [13-22], or as biofertilizer from their N-fixing endosymbionts [for example similar to other species of *Epithemia*, *Ep. turgida* (Bacillariophyceae: Rhopalodiales: Rhopalodiaceae) contains N-fixing Cyanobacteria, these endosymbionts enable algae cells to become abundant in microhabitats with a low N/P ratio and they are frequently abundant as epiphyte on *Cladophora* and other coarse filamentous algae (particularly in western rivers) [23], or as detoxifiers of wastewater polluted with antibiotics (case of *Chaetoceros muelleri* (Bacillariophyta: Bacillariophyceae: Chaetocerotanae *incertae sedis*: Chaetocerotaceae) which is an appealing solution to remove certain antibiotics such as sulfamethoxazole and ofloxacin from wastewater) [24]. Several species of Chromista show a detrimental ecological impact as producers of toxins harmful to aquatic living organisms [25-30]. This is the case of the genus *Ceratium* (Dinophyceae: Gonyaulacales: Ceratiaceae)

which sometimes causes dramatic blooms in lakes and responsible of fish mortality [31]. This is also the case of the genus *Chaetoceros* which is non-toxic to humans but harmful to fish and invertebrates (especially in intensive aquaculture systems) by damaging or clogging their gills [32]. Several species of Chromista are responsible of zoonoses, as the case of oomycetes *Saprolegnia* responsible of the saprolegniosis pathology in fish [33] and the case of labyrinthulomycetes that cause diseases in aquatic animals [34]. Apart from the harmful impact reported in several species, some species (case of labyrinthulomycetes) live as commensals or mutualists within the guts and tissues of aquatic invertebrates and they are saprobic on animal faeces and molluscs shells [34]. These potentialities make Chromists good bio-indicators of the water quality of life.

Nyong and Kienke river mouths (South Cameroon) are source of drinking water and fishing activities [35]. Residents depend on artisanal small-scaled fishing using canoes for household consumption [36, 37] and to supply the neighbouring urban areas [37]. Nevertheless, the demand is growing and fishermen complain about the deterioration of the fish resources for many reasons including irresponsible fishing practices (use of pesticides) and the poor land use management [35]. In this region, the community structure of aquatic micro algae is little known, except works concerning the tidal variation impact on the abundance of phytoplankton in the Nyong estuary [38], seasonal variation of the water quality and the composition of the phytoplankton communities in lower Nyong estuary [39], influence of physico-chemical parameters on the zooplankton dynamics in Kienke estuary [40], the water quality, the biodiversity and abundance of blue-green algae in Nyong and Kienke River mouths [41]. But nothing is known concerning the zoonotic chromists, the toxigenic species, those useful for the nutrition of fish. The place occupied by harmful species in the community. The present study aimed to establish a baseline of information on the distribution of chromists, as a first step in evaluating the status and the occurrence of species known as bio-indicators of the aquatic life quality (useful species or producers of toxins).

## 2. Materials and Methods


### 2.1. Study Sites

Studies took place in 2014 at the Nyong river mouth mouth (03°16'40.71"N, 09°53'27.21"E and 03°14'58.41"N,

09°56'41.07"E) (Figure 1A) and the Kienke river mouth (02°22'4.06"N, 09°48'32.20"E and 02°17'56.31"N, 09°50'55.94"E) (Figure 1B), situated in the Southern coastal zone of Cameroon [41]. They are separated by a distance of 111.1 km. The prevailing climate is tropical with rainfall even during the driest months (December and January: 54.2 mm and 33.8 mm respectively) [41, 42]. The average air temperature ranges from 24.4 °C (August) to 26.7 °C (March) and the average rain fall ranges from 116 mm (January) to 340 mm (September). The average air humidity ranges from 84.0% (January to March) to 87.0% (September and October) [41, 42]. Four seasons are defined: a long dry season (late November-February), a short rainy season (March-June), a short dry season (July-August) and a long rainy season (early September-early November) [36, 41]. Soils are acidic, yellow ferrallitic types, poor in minerals and organic matters and soils on gneiss outcrop cover the bulk between Campo and Kribi [36, 41]. Many streams crossing the region are influenced by the equatorial climate [36, 41]. The main rivers (Nyong, Lokoundje, Kienke, Lobe and Ntem) flow into the Atlantic Ocean and the watercourses are used by the residents for traditional fishing or as waterways using canoes or other navigation fleet [36, 41]. According to our recently published report [41] Nyong and Kienke river mouths belong to the warm category (temperature varying: 26.0 to 32.0 °C). The pH varies from slightly acidic (pH=6.1) to slightly basic (pH=8.9). The transparency varies from 43.0 cm to 325 cm. The dissolved oxygen (DO) varies from 0.4 to 39.5 mg.l<sup>-1</sup>. The five-day biochemical oxygen demand (BOD<sub>5</sub>) varies from 5.0 to 50.0 mg.l<sup>-1</sup>. The conductivity varies from 16.4 to 40,600.0 µS.cm<sup>-1</sup>. The nitrite NO<sub>2</sub><sup>-</sup> contents varies from 0 to 27.0 mg.l<sup>-1</sup>. The nitrate NO<sub>3</sub><sup>-</sup> contents varies from 0 to 9.9 mg.l<sup>-1</sup>. Ammoniacal nitrogen (NH<sub>4</sub><sup>+</sup>) varies from 0.2 to 14.1 mg.l<sup>-1</sup>. Orthophosphate (PO<sub>4</sub><sup>3-</sup>) content varies from 0 to 2.0 mg.l<sup>-1</sup>. The chlorophyll a content varies from 0.02 to 0.40 µg.l<sup>-1</sup>. The biomass varies from 0.3 to 12.0 mg.c.l<sup>-1</sup>. The faecal coliforms content varies from 75.0 to 1440.0 CFU. (100 ml)<sup>-1</sup> and the total suspended solids (TSS) varies from 0 to 33.3 mg.l<sup>-1</sup>.

### 2.2. Sampling Design

Samplings were set up from March to June 2014 in the lower course of Nyong and Kienke mouths (Figure 1A), at the same sampling points presented in our recent publication [41]: four sites were selected 300 m from the shore of Nyong and 30 m from the shore for Kienke.



**Figure 1.** Location of the study sites in Southern coastal zone of Cameroon (southern province, Ocean department). A: Location of the Nyong and Kienke River mouths; B: Location of the collection sites in the Nyong River mouth; C: Location of the collection sites in the Kienke River mouth.

In each river, sampling sites were accessed using a wooden canoe (Nyong River mouth: site 1 at the beginning of the estuary (3°16'1.79"N, 9°56'25.72"E), site 2 at the middle of the estuary: (3°15'57.58"N, 9°55'31.13"E; 1.71 km from site 1), site 3 at the transition with ocean water (3°15'38.99"N, 9°54'16.28"E; 4.11 km and 2.47 km from site 1 and site 2 respectively) and site 4 located in the coastal area of the ocean (3°16'3.85"N, 9°53'45.64"E; 5.15 km, 3.4 km and 1.3 km from site 1, site 2 and site 3 respectively)) (Figure 1B) [41]. In the Kienke River mouth, locations of the sampling sites were: site 1 near a residential camp (2°19'20.40"N,

9°50'17.78"E), site 2 near a marshy area (2°20'14.06"N, 9°50'1.07"E; 1.9 km from site 1), site 3 at the transition with the ocean (2°20'55.01"N, 9°49'22.47"E; 1.8 km and 3.5 km from site 2 and site 1 respectively) and site 4 located in the coastal area of the ocean (2°21'17.29"N, 9°48'57.40"E; 4.8 km, 2.8 km and 1.1 km from site 1, site 2 and site 3 respectively) (Figure 1C) [41]. In each river mouth, as site 4 was situated far from the coast, an outboard boat permitted us to reach it. Samsung 14.2 Mega Pixels and Kodak 9.2 Mega Pixels cameras were used for the field shots. Coordinates of the sites were taken using a Garmin GPS. As presented in our



recent publication [41], three sampling sessions were done at each site (one in March, June and August respectively). The floating macro-particles were sampled and preserved in a water sample in plastic petri dishes hermetically closed with a plastic lid. For the microalgae detection, raw water was sampled at the water surface and one-meter depth using a Teflon ball and a two-liter Niskin messenger bottles and 150 ml transparent plastic polyethylene bottles and transported to the laboratory using a 100 ml Coleman cooler containing pieces of ice for temperature maintenance. Biological parameters of the sampled water (Chromista composition, identification and counting) were carried out at the microalgae laboratory of the Specialized Center for Research on Marine Ecosystems at Kribi (AquaSol service). Chromista cells concentration was determined using the Malassez cells. In case of insufficient homogeneity, the assembly was resumed using a new Malassez cell and the rehomogenized raw water. Chromista cells were identified and for each species, we counted the number of cells in ten randomly selected squared grid areas and the average number ( $n_i$ ) of 10 Malassez's squared grids and the final concentration were recorded as  $c_i = n_i \times 10^5 \text{ cells.ml}^{-1}$ . The total number of cells in volume "v" was estimated as  $n_i = c_i \times v$  and data were compiled in a species matrix database.

### 2.3. Species Identification and Data Analysis

Species identification was made using a Zeiss NR183268 series microscope and by referring to the descriptions, drawings, dimensions and photographs in available dichotomous keys [43-46]. Update name of species and their natural environment were obtained by referring to catalogs and websites available online [11, 47-52]. In each assemblage the weighed mean sensitivity (WWS) of the diatom community was determined and the trophic diatom index (TDI) was calculated. Data are given in terms of absolute and relative frequencies. Two independents percentages were compared using the Fisher's exact test. For the simultaneous comparison of several percentages, the asymptotic p-value or the exact p-value was determined using the independent chi-square test or the Fisher-Freeman-Halton test from StatXact software version 3.1.

Alpha diversity analysis allowed the determination of 15 indexes using PAST 3.05 software: absolute abundance of the  $i^{\text{th}}$  species  $n_i$ , observed sample size  $n$ , relative abundance of the  $i^{\text{th}}$  species  $p_i = (n_i/n) \times 100$ , species richness  $S$ , maximum abundance  $n_1$  or  $n_{\text{max}}$ , Margalef's index  $Mg = (S-1)/\ln(n)$ , richness ratio  $d = S/n$  (0 for the low-rich communities and +1 for the optimal species-rich communities), Shannon-Weaver diversity index  $H'$  (0 for a single-species community and  $H'_{\text{max}} = \ln(S)$  for the perfect species regularity of abundances), Simpson diversity index  $D$  (0 for a high diversities and +1 for a low diversities), Hill's  $N_1$  diversity number ( $N_1 = e^{H'}$ ) for the estimated number of abundant species, Hill's  $N_2$  diversity number ( $N_2 = 1/D$ ) for the estimated number of co-dominants,

Hill's ratio  $N_2/N_1$  with  $0 \leq \text{Hill} \leq +1$ , Pielou's evenness index  $J = H'/\ln(S)$  with 0 for a perfect heterogeneity of the assemblage and +1 for a perfect balance of abundances, and Berger-parker dominance index  $n_{\text{max}}/n$  with a low value reflecting a high diversity. Comparison of the species richness was performed using the individual rarefaction procedure. The non-parametric estimator Chao1 was used to estimate the theoretical species richness  $T$  and the sampling effort was estimated as  $(S/T) \times 100$ . The rank abundance plottings were used to illustrate the shape of the species abundance distributions (SADs). The goodness of fit of each SAD to a theoretical model was assessed by calculating the Pearson correlation between the logarithms of the numbers and the ranks of the species and interpreted as follows:  $r < -0.95$  for a fit of a poor quality;  $r \approx -0.95$  for an approximative fit;  $r \approx -0.98$  for a satisfactory fit; and  $r \geq -0.99$  for an excellent fit. We used five commonly used theoretical models to fit the curves: broken-stick (BS), log-linear (LL) model, lognormal (LN) model, Zipf (Z) and Zipf-Mandelbrot (ZM). The best model was selected using the Akaike Information Criteria (AIC) or the Bayesian Information Criteria (BIC) (the best model presented the lowest AIC or BIC). The estimated sample size  $n^*$  was adjusted to the observed size  $n$  using the correction factor  $n/n^*$  and the corrected models were given. The package *vegan* of R 3.4.1 software helped us to adjust the SADs. BS model (McArthur's model) or model of contiguous non-overlapping niches is based on a hypothetical form of sharing of biotope resources between the species present and in practical, this model is suitable for the analysis of communities in which inter-species relationships are elementary, competition being essentially limited to the level of the resource, such as physical space. This model has a single parameter  $x$  which represents the average abundance of species [53]. LL model (preemption model) suggests that numbers of species are distributed according to the geometrical progression (Motomura's law) and the parameter  $m$  (Motomura's environmental constant with  $0 \leq m \leq 1$ ) is the antilogarithm of the linear slope obtained by plotting species ranks (from the most to the least abundant) on the x-axis and the logarithms of corresponding abundances on the y-axis.

The species of the  $i^{\text{th}}$  rank in the ranking by decreasing abundance has therefore a number of individuals proportional to  $(k)(1-k)^{(i-1)}$ , its logarithm being of the form  $(i-1)\text{Log}(1-k) + \text{Log}(k)$  which is a straight line of slope  $\text{Log}(1-k)$ . This model depends on the maximum abundance of the top-ranking species  $n_1$  and the Motomura's environmental constant  $m$  (the rate of decrease in abundance by rank) [54]. According to the LN model, the logarithms of the abundances are distributed symmetrically and randomly, on either side of their mean, with a standard deviation of the logarithm distribution  $\sigma = \text{square root of } 1/m'$ , with  $m'$  as the Preston's environmental constant. The adjustment line  $\text{Log}(n_i) = f(P_i)$  with  $P_i$  the probit of the cumulative percentage linked to the ranks  $i$ , represents all the points, and makes it possible to calculate the theoretical abundances which would

be those of  $n$  species if the nomocenosis was rigorously log-normal. For a species of rank  $i$ , we calculated the cumulative percentage linked to this rank  $k_i = 100(i+0.5)/(S+1)$  when  $S$  was odd or  $k_i = 100((i+1)+0.5)/(S+1)$  when  $S$  was even, and then the probit  $P_i = P_{(k_i)}$  was determined. The package “ecotoxycology” of R 4.1.0 software helped us to determine the probits. The regression line between  $\text{Log}(n_i)$  and  $P_i$  has the relation:  $\text{Log}(n_i) = aP_i + b$  or  $n_i = (10^b)(10^a)^{P_i}$  where  $a$  and  $b$  represented the slope and the elevation respectively of the regression line  $\text{Log}(n_i) = f(P_i)$ . Z model is based on the Zipf's law [55], with abundances listed in descending order. Z model is defined from two statistics:  $Q$  which is the sum of all recorded abundances, also called sample size or the scale parameter (normalization constant) and  $\gamma$  (gamma) which is the decay coefficient or the average probability of appearance of a species and the relation is  $n_i = Q(i)^{-\gamma}$  where  $i$  represents the rank of the species ranked in decreasing order [56, 57]. ZM model  $n_i = Q(i+\beta)^{-\gamma}$  is a generalized model in which a new parameter  $\beta$  (beta) is added (the degree of the niche diversification). ZM model characterizes evolved ecosystems, where  $n_i$  is the abundance of a particular species of the  $i^{\text{th}}$  rank in the assemblage, the abundances being ranked in decreasing order,  $Q = n_1(1+\beta)^\gamma$  represents the normalizing constant,  $n_1$  is the maximum species abundance,  $S$  is the total number of species and  $1/\gamma$  represents the fractal dimension of the distribution of individuals among species [58, 59]. Marquardt's nonlinear least squares algorithm summarized by Le *et al.* [57] and by Murthy [60], was used to estimate  $\beta$  and  $\gamma$ , iterations starting with an initial guess value  $x_0 = (0; 2)^T$  at  $k=0$ , a tolerance of the functional value  $\epsilon$ , the damping factor  $\lambda$  and the modest value  $\lambda_0 = 100$ . For a vector  $x_i$  we computed the equation  $n_i(1+\beta)^\gamma = Q$ , the gradient  $\nabla f(x_i)$  and the Hessian matrix  $H_i$ . At the  $k$ th iteration  $f(x_k)$  was determined and the equation  $x_{k+1} = x_k - (H + \lambda I)^{-1} \nabla f(x_i)$  was solved and  $f(x_{k+1})$  was determined. When  $f(x_{k+1}) < f(x_k)$  we changed the value of  $\lambda_{k+1} = \lambda_k/2$  and when  $f(x_{k+1}) > f(x_k)$  we changed the value of  $\lambda_{k+1} = 2\lambda_k$ . Iteration was stopped when the solution met the desired convergence criteria  $|f(x_k) - f(x_{k-1})| < \epsilon$ .

For the beta diversity analysis, the dissimilarity between the collection sites, mouths and rivers was evaluated using the Bray-Cutis index. The overall species covariance was evaluated using Schluter's procedure [61]. Between species correlation was evaluated using the Kendall's tau correlation.

## 3. Results

### 3.1. Inventory of Species and Abundances

A total of  $10,427.1 \times 10^5$  cells were collected (dry season:  $833.3 \times 10^5$  cells (8.0%); rainy season:  $9,593.8 \times 10^5$  cells (92.0%)), divided into  $2,458.3 \times 10^5$  cells (23.6%) from Nyong (dry season:  $145.8 \times 10^5$  cells (1.4%); rainy season:  $2,312.5 \times 10^5$  cells (22.2%)) and  $7,968.8 \times 10^5$  cells (76.4%) from Kienke (dry season:  $687.5 \times 10^5$  cells (6.6%); rainy sea-

son:  $7,281.3 \times 10^5$  cells (69.8%)) (Table 1). Cells belonged to three phyla, eight classes, 23 orders, 32 genera and 40 species. Phyla were Dinoflagellata Bütschli 1885 *sensu* Gomez 2012 (33.3% of the total collection) exclusively in Kienke, and two phyla (66.7%) (Bacillariophyta Dillon, 1963, and Ochrophyta Cavalier-Smith, 1995) common to the two mouths (Table 1). Bacillariophyta was the most collected phylum (71.7%), followed by Ochrophyta (22.6.7%), and Dinoflagellata (5.7%). Percentage of Bacillariophyta was significantly high than that of other phyla (Fisher's Exact test with  $df=1$ :  $p=0.004$  in the dry season; Fisher-Freeman-Halton test:  $df=2$ ,  $p<0.001$  in the rainy season and in the pooled rivers respectively). Fisher's exact test showed significantly high occurrence in Kienke than Nyong (Bacillariophyta:  $p=1.2 \times 10^{-41}$  in the dry season;  $p<0.001$  in the rainy season and the pooled rivers; Dinoflagellata:  $p=5.2 \times 10^{-183}$  in the rainy season and the pooled rivers; Ochrophyta:  $p=2.8 \times 10^{-46}$  in the dry season;  $p<0.001$  in the rainy season and the pooled rivers; global assemblage:  $p=5.2 \times 10^{-88}$  in the dry season;  $p<0.001$  in the rainy season and the pooled rivers).

Classes were Xanthophyceae Allorge ex Fritsch, 1935 exclusively in Nyong, three classes (Chrysophyceae Pascher, 1914, Cryptophyceae Fritsch, 1927, and Dinophyceae Fritsch, 1927) exclusively in Kienke, and four classes (Bacillariophyceae Haeckel, 1878, Coscinodiscophyceae Round & R. M. Crawford, 1990, Eustigmatophyceae D. J. Hibberd & Leedale, 1970, and Mediophyceae Medlin & Kaczmarek, 2004) common to the two river mouths. Making five classes (62.5%) in Nyong and seven classes (87.5%) in Kienke. Bacillariophyceae was recorded during all seasons in both mouths. Chrysophyceae and Dinophyceae were recorded only during the rainy season in Kienke. Coscinodiscophyceae was recorded during the rainy season in both mouths. Cryptophyceae was recorded during the dry season in Kienke. Eustigmatophyceae was recorded during the rainy season in both mouths. Mediophyceae was recorded during the dry season in Nyong and during both seasons in Kienke. Xanthophyceae was recorded during the dry season in Nyong. Ochrophyta was the most represented phylum with six classes (75.0%) while Bacillariophyta and Dinoflagellata were represented each by one class (12.5%). Bacillariophyceae was the most recorded class (68.9% of the total collection), followed very far by Coscinodiscophyceae (9.2%), Eustigmatophyceae (7.2%), Dinophyceae (5.7%), Mediophyceae (4.4%), Cryptophyceae (3.6%), while other classes were represented each by less than 1.0% of the total collection. Percentages of classes were significantly higher during the rainy season than the dry season (Fisher's exact test:  $p<0.001$ ) and they significantly were higher in Kienke than in Nyong (Fisher's exact test:  $p<0.001$ ).

Six orders (26.1%) were recorded exclusively in Nyong (Achnanthales P. C. Silva, 1962, Aulacoseirales R. M. Crawford, 1990, Coscinodisciales Round (d) & R. M. Crawford, 1990, Mischococcales F. E. Fritsch, 1927 (Xantho-

phyceae), Rhabdonematales Round (d) & R. M. Crawford (d), 1990, and Thalassiophysales D. G. Mann, 1990). Ten orders (43.5%) were collected exclusively in Kienke (*Chaetocerotanae incertae sedis* as temporary name since the classification is controversial, Chromulinales Pascher, 1910, Cryptomonadales Pringsheim (d), 1944, Gonyaulacales F. J. R. Taylor, 1980, Mastogloiales D. G. Mann (d), 1990, Melosirales R. M. Crawford (d), 1990, Phytodinales T. Christensen, 1962 ex Loeblich 1970, Rhizosoleniales P. C. Silva (d), 1962, Rhopalodiales D. G. Mann, 1990, and Thalassiosirales Glezer (d) & I. V. Makarova (d), 1986). Sevent orders (30.4%) were common to the two river mouths (Bacillariales Hendey, 1937 *sensu emend*, Cymbellales D. G. Mann, 1990, Fragilariales P. C. Silva (d), 1962, Goniochloridales K. P. Fawley (d), M. Eliáš (d) & M. W. Fawley (d), 2014, Naviculales Bessey, 1907, Stephanodiscals Nikolaev (d) & Harwood (d), 1997, and Surirellales D. G. Mann (d), 1990). Making 13 orders (56.5%) in Nyong and 17 orders (73.9%) in Kienke. The most represented class was Bacillariophyceae (52.2% of the recorded orders; 68.9% of the total collection), followed by Coscinodiscophyceae (17.4% of the recorded orders and 9.2% of the total collection), Dinophyceae (8.7% of the recorded orders and 5.7% of the total collection), Eustigmatophyceae (4.3% of the recorded orders and 7.2% of the total collection), Mediophyceae (4.3% of the recorded orders and 4.4% of the total collection), Cryptophyceae (4.3% of the recorded orders and 3.6% of the total collection), while Chrysophyceae and Xanthophyceae were each represented by one order (4.3% of the recorded orders, and less than 1.0% of the total collection). Melosirales was the most collected order (13.4% of the total collection), followed by Bacillariales (12.2%), Cymbellales (10.8%), Surirellales (10.8%), Goniochloridales (7.2%), Rhizosoleniales (7.0%), Fragilariales (6.3%), Thalassiosirales (3.9%), Stephanodiscals (3.8%), Cryptomonadales (3.6%), Mischococcales (3.6%), Cryptomonadales (3.5%), Naviculales (3.2%), Gonyaulacales (3.0%), Thalassiophysales (2.8%), Phytodinales (2.7%), Achnanthales (2.6%), Aulacoseirales (2.2%), and Rhopalodiales (1.2%). Other orders were represented each by less than 1.0% of the total collection.

Ten families (38.5%) were recorded exclusively in Nyong (Achnanthaceae, Amphipleuraceae, Anomoeoneidaceae, Aulacoseiraceae, Catenulaceae, Cocconeidaceae, Coscinodiscaceae, Ophiocytaceae, Pinnulariaceae, and Tabellariaceae) (Table 1). Ten families (38.5%) were recorded exclusively in Kienke (Ceratiaceae, Chaetocerotaceae, Cryptomonadaceae, Diploneidaceae, Gomphonemataceae, Mastogloaceae, Melosiraceae, Phytodiniaceae, Rhizosoleniaceae, and Rhopalodiaceae) (Table 1). Six families (23.1%) were common to both river mouths (Bacillariaceae, Dinobryaceae, Fragilariaceae, Goniochloridaceae, Stephanodiscaceae, and Surirellaceae) (Table 1). Making 16 families (61.5%) in Nyong and Kienke respectively. Based on the number of families, the most represented order was Naviculales [three families (11.5%, 3.1% of the total collection) (Table 1). It

was followed by Achnanthales and Cymbellales [two families each (7.7%), 2.6% and 10.8% of the total collection respectively] (Table 1). Other orders were each represented by one family (3.8%) (Table 1). Melosiraceae was the most collected family (13.4% of the total collection), followed by Bacillariaceae (12.2%), Surirellaceae (10.8%), Gomphonemataceae (9.6%), Stephanodiscaceae (7.7%), Goniochloridaceae (7.2%), Rhizosoleniaceae (7.0%), Fragilariaceae (6.3%), Cryptomonadaceae (3.5%), Ceratiaceae (3.0%), Catenulaceae (2.8%), Phytodiniaceae (2.6%), Aulacoseiraceae (2.2%), Diploneidaceae (1.9%), Cocconeidaceae (1.4%), Achnanthaceae (1.2%), Anomoeoneidaceae (1.2%), Rhopalodiaceae (1.2%) (Table 1). Other families were rare and represented each by less than 1.0% of the total collection. Twenty species (50.0%) and no common species were recorded in each river mouth (Table 1). Five species (12.5%) were collected in the dry season (Table 1). Fifteen species (37.5%) were collected in the rainy season (Table 1).

Twenty-one species (52.5%) were common to both seasons (Table 1). In the pooled data from the two river mouths, six species (15.0%) were collected in the dry season and 34 species (85.0%) were collected in the rainy season (Table 1).

In Nyong, three species (7.5%) were recorded in the dry season (*Cyclotella meneghiniana* var. *meneghiniana* Kützing, 1844 (Stephanodiscaceae), *Denticula elegans* Kützing, 1844 (Bacillariaceae), and *Ophiocytium cochleare* (Eichwald) A. Braun 1855 (Ophiocytaceae)) (Table 1). Seventeen species (42.5%) were collected exclusively in the rainy season (*Achnanthes exiguides* Compère, 1967 (Achnanthaceae), *Amphora ovalis* (Kützing) Kützing, 1844 (Catenulaceae), *Anomoeoneis sphaerophora* E. Pfitzer, 1871 (Anomoeoneidaceae), *Aulacoseira granulata* (Ehrenberg) Simonsen, 1979 (Aulacoseiraceae), *Campylodiscus noricus* Ehrenberg ex Kützing, 1844 (Surirellaceae), *Cocconeis placentula* var. *placentula* Ehrenberg 1838 *sensu* Jahn et al. 2009 (Coscinodiscaceae), *Coscinodiscus rudolfi* Bachmann, 1938 (Coscinodiscaceae), *Cymatopleura solea* var. *apiculata* W. Smith, 1853 (Surirellaceae), *Cymatopleura solea* var. *bai-calensis* Skvortzow & Meyer, 1928 (Surirellaceae), *Diploneis arctica* (Lange-Bertalot) Lange-Bertalot & A. Fuhrmann, 2016 (Diploneidaceae), *Fragilaria construens* f. *construens* (Ehrenberg) Grunow, 1862 (Fragilariaceae), *Frustulia adnata* Kützing, 1833 (Amphipleuraceae), *Goniochloris mutica* (A. Braun) Fott, 1960 (Goniochloridaceae), *Hantzschia amphioxys* (Ehrenberg) Grunow, 1880 (Bacillariaceae), *Nitzschia sigma* (Kützing) W. Smith, 1853 (Bacillariaceae), *Pinnularia cardinaliculus* var. *ceylonica* Skvortzow, 1930 (Pinnulariaceae), and *Tabellaria flocculosa* (Roth) Kützing, 1844 (Tabellariaceae)) (Table 1). No common species was recorded.

In Kienke, three species (7.5%) were recorded exclusively in dry season (*Chaetoceros muelleri* Lemmermann, 1898 (Chaetocerotaceae), *Cryptomonas erosa* Ehrenberg 1832 (Cryptomonadaceae), and *Surirella linearis* f. *kolhapurensis* Sarode & Kamat, 1984 (Surirellaceae)) (Table 1).

Seventeen species (42.5%) were collected exclusively during the rainy season (*Ceratium hirundinella* (O. F. Müller) Dujardin, 1841 (Ceratidae), *Cyclotella stelligera* Cleve & Grunow, 1882 (Stephanodiscaceae), *Denticula thermalis* var. *fossilis* Frenguelli, 1936 (Bacillariaceae), *Diploneis ovalis* var. *pumila* (Grunow) Cleve, 1894 (Diploneidaceae), *Cystodinium unicorne* G. A. Klebs, 1912 (Phytodiniaceae), *Dinobryon sertularia* Ehrenberg, 1834 (Dinobryaceae), *Epithemia turgida* var. *turgida* (Ehrenberg) Kützing, 1844 (Rhopalodiaceae), *Gomphonema olivaceum* (Lyngbye) Desmazières, 1825 (Gomphonemataceae), *Goniochloris gigas* Bourrelly (Goniochloridaceae), *Mastogloia smithii* Thwaites ex W. Smith, 1856 (Mastogloiaceae), *Nitzschia amphibia* Grunow, 1862 (Bacillariaceae)) (Table 1). Other species were *Rhizosolenia longiseta* O. Zacharias, 1893 (Rhizosoleniaceae), *Stephanodiscus astra* (Ehrenberg) Grunow, 1880 (Stephanodiscaceae), *Surirella capronii* Brébisson & Kitton, 1869 (Surirellaceae), and *Synedra ulna*

(Nitzsch) Ehrenberg, 1832 (Fragilariaceae) (Table 1). No species was common to both seasons. In the pooled distribution, the most specious family was Bacillariaceae [six species (15.0%): three species (7.5%) in each river mouth], followed by Surirellaceae (five species (12.5%); two species (5.0%) in Kienke and three species (7.5%) in Nyong), Stephanodiscaceae (three species (7.5%); one species (2.5%) in Nyong and two species (5.0%) in Kienke) (Table 1). Diploneidaceae and Fragilariaceae were each represented by two species (5.0% divided into one species (2.5%) in Nyong and Kienke respectively). Other families were each represented by one species (2.5%) in Nyong or Kienke (Table 1). *Me. granulata* was the most collected species (13.4%), followed by *Go. olivaceum* (9.6%), *Rh. longiseta* (7.0%), *Gn. gigas* (6.4%), *De. thermalis* and *Sy. ulna* (5.1% respectively), *St. astra* (3.9%), *Ca. noricus* (3.7%), *Cr. erosa* (3.6%), *Ni. amphibia* (3.6%), *Cc. stelligera* (3.2%), *Ce. hirundinella* (3.0%), *Am. ovalis* (2.8%), *Cy. unicorne* (2.6%), *Su. linearis* (2.4%), *Su. capronii* (2.3%), *Au. granulata* (2.2%), *Ha. amphioxys* (2.1%). Other species were rare (Table 1).

**Table 1.** Absolute and relative abundances of the Chromista species in the Nyong (A) and Kienke (B) River mouths.

Families/Species	References	A1 (%)	A2 (%)	Total (%)	B1 (%)	B2 (%)	Total (%)
Achnanthaceae Kütz., 1844							
<i>Ac. exiguides</i> <sup>#,BI</sup>	[43]	-	125.0 (1.2)	125.0 (1.2)	-	-	-
Amphileuraceae Grunow (d), 1862							
<i>Fr. adnata</i> <sup>#,BI</sup>	[49]	-	62.5 (0.6)	62.5 (0.6)	-	-	-
Anomoeoneidaceae D. G. Mann, 1990							
<i>An. sphaerophora</i> <sup>*,#,BI</sup>	[21, 43]	-	125.0 (1.2)	125.0 (1.2)	-	-	-
Aulacoseiraceae R. M. Crawford, 1990							
<i>Au. granulata</i> <sup>#,EI</sup>	[13, 19-21, 48, 49]	-	229.2 (2.2)	229.2 (2.2)	-	-	-
Bacillariaceae Ehrenb., 1831							
<i>De. elegans</i> <sup>#,BI</sup>	[11]	62.5 (0.6)	-	62.5 (0.6)	-	-	-
<i>De. thermalis</i> <sup>*,BI</sup>	[11]	-	-	-	-	531.3 (5.1)	531.3 (5.1)
<i>Ha. amphioxys</i> <sup>†,#,*,BI</sup>	[21, 43, 49]	-	218.8 (2.1)	218.8 (2.1)	-	-	-
<i>Ni. amphibia</i> <sup>#,BI</sup>	[16, 22, 43, 48, 49]	-	-	-	-	375.0 (3.6)	375.0 (3.6)
<i>Ni. sigma</i> <sup>#,BI</sup>	[22, 43, 49]	-	20.8 (0.2)	20.8 (0.2)	-	-	-
<i>Ni. tryblionella</i> <sup>#,†,BI</sup>	[43]	-	-	-	-	62.5 (0.6)	62.5 (0.6)
Catenulaceae Mereschkowsky, 1902							
<i>Amphora ovalis</i> <sup>#,BI</sup>	[21, 43]	-	291.7 (2.8)	291.7 (2.8)	-	-	-
Ceratidae Kofoid, 1907							
<i>Ce. hirundinella</i> <sup>#,†,BL</sup>	[44]	-	-	-	-	312.5 (3.0)	312.5 (3.0)



Families/Species	References	A1 (%)	A2 (%)	Total (%)	B1 (%)	B2 (%)	Total (%)
Chaetocerotaceae Ralfs (d), 1861							
<i>Ch. muelleri</i> <sup>†,US,ITP</sup>	[21]	-	-	-	62.5 (0.6)	-	62.5 (0.6)
Cocconeidaceae Kützing, 1844							
<i>Co. placentula</i> <sup>#,†,BI</sup>	[14, 21, 43, 48]	-	145.8 (1.4)	145.8 (1.4)	-	-	-
Coscinodiscaceae Kützing, 1844							
<i>Cs. rudolfi</i> <sup>†,BI</sup>	[15, 43]	-	62.5 (0.6)	62.5 (0.6)	-	-	-
Cryptomonadaceae Ehrenberg, 1831							
<i>Cr. erosa</i> <sup>*,#,BI</sup>	[52]	-	-	-	375.0 (3.5)	-	375.0 (3.5)
Dinobryaceae Ehrenberg, 1834							
<i>Di. sertularia</i> <sup>*,#,BI</sup>	[52]	-	-	-	-	83.3 (0.8)	83.3 (0.8)
Diploneidaceae D. G. Mann (d), 1990							
<i>Dp. arctica</i> <sup>#,BI</sup>	[63]	-	62.5 (0.6)	62.5 (0.6)	-	-	-
<i>Dp. ovalis</i> <sup>#,BI</sup>	[11]	-	-	-	-	145.8 (1.4)	145.8 (1.4)
Fragilariaceae Kützing, 1844							
<i>Fr. construens</i> <sup>*,#,BI</sup>	[13, 21, 43]	-	125.0 (1.2)	125.0 (1.2)	-	-	-
<i>Synedra ulna</i> <sup>#,BI</sup>	[16, 43, 49]	-	-	-	-	531.3 (5.1)	531.3 (5.1)
Gomphonemataceae Kützing, 1844							
<i>Go. olivaceum</i> <sup>#,BI</sup>	[16, 48]	-	-	-	-	1000.0 (9.6)	1000.0 (9.6)
Goniochloridaceae Bailey (d)							
<i>Gn. gigas</i> <sup>#,BI</sup>	[43]	-	-	-	-	666.7 (6.4)	666.7 (6.4)
<i>Gn. mutica</i> <sup>#,BI</sup>	[45]	-	83.3 (0.8)	83.3 (0.8)	-	-	-
Mastogloiaceae Mereschkowsky, 1903							
<i>Ma. smithii</i> <sup>#,BI</sup>	[43]	-	-	-	-	62.5 (0.6)	62.5 (0.6)
Melosiraceae Kützing, 1844							
<i>Me. granulata</i> <sup>#,†,BI</sup>	[43, 48]	-	-	-	-	1395.8 (13.4)	1395.8 (13.4)
Ophiocytaceae Lemmermann, 1899							
<i>Op. cochleare</i> <sup>#,BI</sup>	[50]	20.8 (0.2)	-	20.8 (0.2)	-	-	-
Phytodiniaceae Klebs, 1912							
<i>Cy. unicorn</i> <sup>#,BI</sup>	[44]	-	-	-	-	281.3 (2.6)	281.3 (2.6)
Pinnulariaceae D. G. Mann (d), 1990							
<i>Pi. cardinaliculus</i> <sup>#,BI</sup>	[62]	-	62.5 (0.6)	62.5 (0.6)	-	-	-
Rhizosoleniaceae De Toni, 1890							
<i>Rh. longiseta</i> <sup>#,BI</sup>	[43]	-	-	-	-	729.2 (7.0)	729.2 (7.0)
Rhopalodiaceae Topachevs'kyj (d) & Oksiyuk (d), 1960							
<i>Ep. turgida</i> <sup>†,US(NF)</sup>	[16, 23, 48]	-	-	-	-	125.0 (1.2)	125.0 (1.2)
Stephanodiscaceae I. V. Makarova (d), 1986							
<i>Cc. meneghiniana</i> <sup>#,†,BI</sup>	[43, 48, 49]	62.5 (0.6)	-	62.5 (0.6)	-	-	-

Families/Species	References	A1 (%)	A2 (%)	Total (%)	B1 (%)	B2 (%)	Total (%)
<i>Cc. stelligera</i> <sup>#,BI</sup>	[17, 43, 48, 49]	-	-	-	-	333.3 (3.2)	333.3 (3.2)
<i>St. astraea</i> <sup>#,†,BI</sup>	[13]	-	-	-	-	406.3(3.9)	406.3(3.9)
Suriellaceae Kützinger, 1844							
<i>Ca. noricus</i> <sup>#,BI</sup>	[11]	-	385.4 (3.7)	385.4 (3.7)	-	-	-
<i>Cm. apiculata</i> <sup>#,BI</sup>	[21]	-	62.5 (0.6)	62.5 (0.6)	-	-	-
<i>Cm. solea</i> <sup>#,BI</sup>	[21]	-	187.5 (1.8)	187.5 (1.8)	-	-	-
<i>Su. capronii</i> <sup>#,BI</sup>	[43]	-	-	-	-	239.6 (2.3)	239.6 (2.3)
<i>Su. linearis</i> <sup>#,BI</sup>	[43]	-	-	-	250.0(2.3)	-	250.0 (2.3)
Tabellariaceae Kützinger, 1844							
<i>Ta. flocculosa</i> <sup>#,‡,BI</sup>	[13, 22, 43, 48, 49]	-	62.5 (0.6)	62.5 (0.6)	-	-	-
Total		145.8(1.4)	2312.5(22.2)	2458.3(23.6)	687.5(6.6)	7281.3(69.8)	7968.8(76.4)

Table 1. Continued.

Families	Species	References	C1 (%)	C2 (%)	Total (%)
Achnanthaceae	<i>Achnanthes exiguides</i> <sup>#,BI</sup>	[43]	-	125.0 (1.2)	125.0 (1.2)
Amphipleuraceae	<i>Frustulia adnata</i> <sup>#,BI</sup>	[49]	-	62.5 (0.6)	62.5 (0.6)
Anomoeoneidaceae	<i>Anomoeoneis sphaerophora</i> <sup>*,#,BI</sup>	[21]	-	125.0 (1.2)	125.0 (1.2)
Aulacoseiraceae	<i>Aulacoseira granulata</i> <sup>#,EI</sup>	[13, 19-21, 48, 49]	-	229.2 (2.2)	229.2 (2.2)
Bacillariaceae	<i>Denticula elegans</i> <sup>#,BI</sup>	[11]	62.5 (0.6)	-	62.5 (0.6)
	<i>De. thermalis</i> var. <i>fossilis</i> <sup>*,BI</sup>	[11]	-	531.3 (5.1)	531.3 (5.1)
	<i>Hantzschia amphioxys</i> <sup>†,‡,BI</sup>	[21, 43, 49]	-	218.8 (2.1)	218.8 (2.1)
	<i>Nitzschia amphibia</i> <sup>#,BI</sup>	[16, 18, 21, 43, 48, 49]	-	375.0 (3.6)	375.0 (3.6)
	<i>Ni. sigma</i> <sup>#,BI</sup>	[21, 43, 49]	-	20.8 (0.2)	20.8 (0.2)
	<i>Ni. tryblionella</i> <sup>#,†,BI</sup>	[43]	-	62.5 (0.6)	62.5 (0.6)
Catenulaceae	<i>Amphora ovalis</i> <sup>#,BI</sup>	[21, 43]	-	291.7 (2.8)	291.7 (2.8)
Ceratiaceae	<i>Ceratium hirundinella</i> <sup>#,†,BI</sup>	[44]	-	312.5 (3.0)	312.5 (3.0)
Chaetocerotaceae	<i>Chaetoceros muelleri</i> <sup>†,ITP</sup>	[21]	62.5 (0.6)	-	62.5 (0.6)
Cocconeidaceae	<i>Cocconeis placentula</i> <sup>#,†,BI</sup>	[14, 21, 43, 48]	-	145.8 (1.4)	145.8 (1.4)
Coscinodiscaceae	<i>Coscinodiscus rudolfi</i> <sup>†,BI</sup>	[15, 43]	-	62.5 (0.6)	62.5 (0.6)
Cryptomonadaceae	<i>Cryptomonas erosa</i> <sup>*,#,BI</sup>	[52]	375.0 (3.6)	-	375.0 (3.6)
Dinobryaceae	<i>Dinobryon sertularia</i> <sup>*,#,BI</sup>	[52]	-	83.3 (0.8)	83.3 (0.8)
Diploneidaceae	<i>Diploneis arctica</i> <sup>#,BI</sup>	[63]	-	62.5 (0.6)	62.5 (0.6)
	<i>Dp. ovalis</i> var. <i>pumila</i> <sup>#,BI</sup>	[11]	-	145.8 (1.4)	145.8 (1.4)
Fragilariaceae	<i>Fragilaria construens</i> <sup>*,#,BI</sup>	[13, 16, 21, 43]	-	125.0 (1.2)	125.0 (1.2)
	<i>Synedra ulna</i> <sup>#,BI</sup>	[43, 49]	-	531.3 (5.1)	531.3 (5.1)
Gomphonemataceae	<i>Gomphonema olivaceum</i> <sup>#,BI</sup>	[16, 48]	-	1000.0 (9.6)	1000.0 (9.6)

Families	Species	References	C1 (%)	C2 (%)	Total (%)
Goniocladoriaceae	<i>Goniocloris gigas</i> <sup>#,BI</sup>	[43]	-	666.7 (6.4)	666.7 (6.4)
	<i>Gn. mutica</i> <sup>#,UN(BI)</sup>	[45]	-	83.3 (0.8)	83.3 (0.8)
Mastogloiaaceae	<i>Mastogloia smithii</i> <sup>#,BI</sup>	[43]	-	62.5 (0.6)	62.5 (0.6)
Melosiraceae	<i>Melosira granulata</i> <sup>#,†,BI</sup>	[43, 48]	-	1395.8(13.4)	1395.8 (13.4)
Ophiocytaceae	<i>Ophiocytium cochleare</i> <sup>#,BI</sup>	[50]	20.8 (0.2)	-	20.8 (0.2)
Phytodiniaceae	<i>Cystodinium unicolor</i> <sup>#,BI</sup>	[44]	-	281.3 (2.6)	281.3 (2.6)
Pinnulariaceae	<i>Pinnularia cardinaliculus</i> <sup>#,BI</sup>	[62]	-	62.5 (0.6)	62.5 (0.6)
Rhizosoleniaceae	<i>Rhizosolenia longiseta</i> <sup>#,BI</sup>	[43]	-	729.2 (7.0)	729.2 (7.0)
Rhopalodiaceae	<i>Epithemia turgida</i> <sup>†, US(BF)</sup>	[16, 23, 48]	-	125.0 (1.2)	125.0 (1.2)
Stephanodiscaceae	<i>Cyclotella meneghiniana</i> <sup>#,†,BI</sup>	[21, 43, 48, 49]	62.5 (0.6)	-	62.5 (0.6)
	<i>Cc. stelligera</i> <sup>#,BI</sup>	[17, 43, 48, 49]	-	333.3 (3.2)	333.3 (3.2)
	<i>Stephanodiscus astraes</i> <sup>#,†,BI</sup>	[13]	-	406.3 (3.9)	406.3 (3.9)
Surirellaceae	<i>Campylodiscus noricus</i> <sup>#,BI</sup>	[11]	-	385.4 (3.7)	385.4 (3.7)
	<i>Cm. apiculata</i> <sup>#,BI</sup>	[21]	-	62.5 (0.6)	62.5 (0.6)
	<i>Cm. solea</i> <sup>#,BI</sup>	[21]	-	187.5 (1.8)	187.5 (1.8)
	<i>Surirella capronii</i> <sup>#,BI</sup>	[43]	-	239.6 (2.3)	239.6 (2.3)
	<i>Su. linearis</i> <sup>#,BI</sup>	[43]	250.0 (2.4)	-	250.0 (2.4)
Tabellariaceae	<i>Ta. flocculosa</i> <sup>#,*,BI</sup>	[13, 16, 21, 22, 43, 48, 49]	-	62.5 (0.6)	62.5 (0.6)
Total			833.3 8.0)	9593.8 92.0)	10427.1(100.0)

A1. Nyong River mouth ( $\times 10^5$  cells) in the dry season; A2. Nyong River mouth ( $\times 10^5$  cells) in the rainy season; B1. Kienke River mouth ( $\times 10^5$  cells) in the dry season; B2. Kienke River mouth ( $\times 10^5$  cells) in the rainy season; C1. Pooled rivers ( $\times 10^5$  cells) in the dry season; C2. Pooled rivers ( $\times 10^5$  cells) in the rainy season; \*: brackish water species; #: freshwater species; †: marine species; ‡: terrestrial species; BF: bio-fertilizer, BI: bio-indicator; BL: bloom forming species; EI: eco-pollution indicator; ITP: ichthyotoxin producer, TS: toxigenic species, WD: widely distributed species.

Based on the water type, 24 freshwater species (60.0%) were divided into 12 species (30.0%) (*Ac. exiguides*, *Am. ovalis*, *Au. granulata*, *Ca. noricus*, *Cm. apiculata*, *Cm. solea*, *De. elegans*, *Dp. arctica*, *Fr. adnata*, *Gn. mutica*, *Ni. sigma*, and *Pi. cardinaliculus*) in Nyong and 12 species (30.0%) (*Cc. stelligera*, *Cy. unicolor*, *Dp. ovalis*, *Gn. gigas*, *Go. olivaceum*, *Ma. smithii*, *Ni. amphibian*, *Op. cochleare*, *Rh. longiseta*, *Su. capronii*, *Su. linearis*, and *Sy. ulna*) in Kienke. Three marine species (7.5%) were recorded (*Cs. rudolfi* in Nyong and two species (*Ch. muelleri*, and *Ep. turgida*) in Kienke). One brackish water specialist (2.5%) (*De. thermalis*) was recorded in Kienke. Thirteen tolerant species (3.5%) were recorded (seven species (17.5%) (*An. sphaerophora*, *Cc. meneghiniana*, *Co. placentula*, *Fa. construens*, *Ha. amphioxys*, *Op. cochleare*, and *Ta. flocculosa*) in Nyong and six species (15.0%) (*Ce. hirundinella*, *Cr. erosa*, *Di. sertularia*, *Me. granulata*, *Ni. tryblionella*, and *St. astraes*) in Kienke) (Table 2). Tolerant species were able to develop in several environments. Seven species (17.5%) could develop in marine water and freshwater (*Cc. meneghiniana*, *Ce. hirundinella*, *Co. placentula*, *Me.*

*granulata*, *Ni. tryblionella*, *Op. cochleare*, and *St. astraes*) (Table 2).

Four species (10.0%) were able to develop in brackish water and freshwater (*An. sphaerophora*, *Cr. erosa*, *Di. sertularia*, and *Fa. construens*) (Table 2). *Ta. flocculosa* was able to develop in freshwater and the terrestrial environments (Table 2). *Ha. amphioxys* was able to develop in the marine, freshwater and terrestrial environments (Table 2). Between habitat specialists (restricted to a single type of water environment) the variation in the percentages was significant except the case of tolerant species in the dry season in Nyong (Table 2). In each river mouth and in each season, among specialist species, freshwater species were significantly more numerous than other categories (Table 2). Freshwater species and/or brackish water were statistically more numerous than species of other categories, except in the dry season in Nyong (Table 2). Between the two rivers, the abundances in Kienke were higher than those recorded in Nyong (Table 2). In each river mouth, the data collected in the rainy season were higher than those recorded in the dry season, both for specialist species and for

tolerant species (Table 2). Specialists statistically outnumber tolerant species, except in the dry season in Nyong (Table 2).

Based on the tides, in Nyong, two species (5.0%) (*De. elegans*, and *Op. cochleare*) were recorded exclusively at high tide during the dry season. One species (2.5%) (*Cc. meneghiniana*) was recorded exclusively at low tide during the dry season. Nine species (22.5%) (*Ac. exiguides*, *An. sphaerophora*, *Cm. apiculata*, *Co. placentula*, *Cs. Rudolphi*, *Dp. arctica*, *Fa. construens*, *Fr. adnata*, and *Ha. amphioxys*) were seen exclusively at high tide in the rainy season. Eight species (20.0%) (*Am. ovalis*, *Au. granulata*, *Ca. noricus*, *Cm. solea*, *Gn. mutica*, *Ni. sigma*, *Pi. cardinaliculus*, and *Ta. flocculosa*) were exclusively at low tide in the rainy season. No species was common to both tides. Making three species (7.5%) in the

dry season and 17 species (42.5%) in the rainy season. In Kienke, one species (2.5%) (*Cr. erosa*) was recorded exclusively at high tide during the dry season. Two species (5.0%) (*Ch. muelleri*, and *Su. linearis*) were recorded exclusively at low tide during the dry season. Two species (5.0%) (*Ma. smithii*, and *Sy. ulna*) were recorded exclusively at high tide during the rainy season. Fourteen species (35.0%) (*Cc. stelligera*, *Ce. hirundinella*, *Cy. unicorn*, *De. thermalis*, *Di. sertularia*, *Dp. ovalis*, *Gn. gigas*, *Go. olivaceum*, *Ep. turgida*, *Me. granulata*, *Ni. amphibia*, *Ni. tryblionella*, *Rh. longiseta*, *St. astraea*, and *Su. capronii*) were collected exclusively at low tide during the rainy season. No species was common to both tides.

**Table 2.** Absolute and relative abundances of the Chromista in three types of water environments.

Water	Nyong River mouth			Kienke River mouth		
	A. (%)	B (%)	Pooled (%)	A (%)	B (%)	Pooled rivers (%)
Abundance of the specialist species x10 <sup>5</sup> (%)						
Freshwater	62.5 (0.6)	1572.9 (15.1)	1635.4 (15.7)	250.0 (2.4)	4364.6 (41.9)	4614.6 (44.3)
Brackish	-	-	-	-	531.3 (5.1)	531.3 (5.1)
Marine	-	62.5 (0.6)	62.5 (0.6)	62.5 (0.6)	125.0 (1.2)	187.5 (1.8)
Total 1	62.5 (0.6)	1635.4 (15.7)	1697.9 (16.3)	312.5 (3.0)	5060.8 (48.5)	5333.3 (51.1)
Test (FE or FFH):	-	p<0.001*	p<0.001*	p<0.001*	FFH: df=2, p<0.001 *	FFH: df=2, p<0.001 *
Avs.B: FE (df=1)	I: p<0.001*; III: p=2.0x10 <sup>-19</sup> *; Total: p<0.001 *			I: p<0.001*; II: p<0.001*; III: p=6.6x10 <sup>-6</sup> *; Total: FE: p<0.001 *		
Abundance of the tolerant species x10 <sup>5</sup> (%)						
I	83.3 (0.8)	145.8 (1.4)	229.2 (2.2)	-	2177.1(20.9)	2177.1 (20.9)
II	-	-	-	375.0 (3.6)	83.3 (0.8)	458.3 (4.4)
III	-	62.5 (0.6)	62.5 (0.6)	-	-	-
IV	-	250.0 (2.4)	250.0 (2.4)	-	-	-
V	-	218.8 (2.1)	218.8 (2.1)	-	-	-
Total 2	83.3 (0.8)	677.1 (6.5)	760.4 (7.3)	375.0 (3.6)	2260.4 (21.7)	2635.4 (25.3)
Global.	145.8 (1.4)	2312.5 (22.2)	2458.3 (23.6)	687.5 (6.6)	7281.3 (69.8)	7968.8 (76.4)
Total 1vs.Total 2	p=0.114 ns	p=.5x10 <sup>-101</sup> *	p<0.001*	p=0.018 *	p<0.001*	p<0.001*
Test	-	FFH: df=3, p<0.001 *	FFH: df=3 p<0.001 *	-	FE: df=1, p<0.001 *	FE: df=1, p<0.001 *
Total 1 vs. Total 2	p<0.001*	p<0.001*	p<0.001*	p=0.018 ns	p<0.001*	p<0.001*
Avs.B (FE)	IV: df=1, p<0.001*; VI: df=1, p<0.001* VII: df=1, p<0.001*; VIII: df=1, p<0.001* Total: df=1, p<0.001*			IV: df=1, p<0.001*; V: df=1, p<0.001*; Total: df=1, p<0.001*		



Table 2. Continued.

Water	Both River mouths		
	A. (%)	B (%)	Pooled rivers (%)
Abundance of the specialist species x10 <sup>5</sup> (%)			
Freshwater	312.5 (3.0)	5937.5 (56.9)	6250.0 (59.9)
Brackish	-	531.3 (5.1)	531.3 (5.1)
Marine	62.5 (0.6)	187.5 (1.8)	250.0 (2.4)
Total 1	375.0 (3.6)	6656.3 (63.8)	7031.3 (67.4)
Test:	FE: df=1, p<0.001 *	FFH: df=2, p<0.001 *	FFH: df=2, p<0.001 *
Avs.B: FE (df=1)	I: p<0.001*; II. p<0.001*; III; p=6.6x10 <sup>-6</sup> *; Total: FE: p<0.001*		
Abundance of the tolerant species x10 <sup>5</sup> (%)			
I	83.3 (0.8)	2322.9 (22.3)	2406.3(23.1)
II	375 (3.6)	83.3 (0.8)	458.3 (4.4)
III	-	62.5 (0.6)	62.5 (0.6)
IV	-	250.0 (2.4)	250.0 (2.4)
V	-	218.8 (2.1)	218.8 (2.1)
Total 2	458.3 (4.4)	2937.5 (28.2)	3395.8 (32.6)
Global.	833.3 (8.0)	9593.8 (92.0)	10427.1 (100.0)
Test	FE: df=1, p<0.001*	FFH: df=4, p<0.001 *	FFH: df=4, p<0.001 *
Avs.B (FE)	IV: df=1, p<0.001*; V: df=1, p<0.001*; VI: df=1, p<0.001*; VII: df=1, p<0.001*; VIII: df=1, p<0.001*; Total: df=1, p<0.001*		
Comparison: Nyong River mouth vs. Kienke River mouth (Fisher’s exact test)			
Freshwater (df=1)	p<0.001*	p<0.001*	p<0.001*
Brackish	-	p<0.001*	p<0.001*
Marine	p<0.001*	p=6.6x10 <sup>-6</sup> *	p=8.6x10 <sup>-16</sup> *
Total 1	p<0.001*	p<0.001*	p<0.001*
I	p<0.001*	p<0.001*	p<0.001*
II	-	p<0.001*	p<0.001*
III	-	p<0.001*	p<0.001*
IV	-	p<0.001*	p<0.001*
V	-	p<0.001*	p<0.001*
Total 2	p<0.001*	p<0.001*	p<0.001*
Global	p<0.001*	p<0.001*	p<0.001*
Total 1 vs. Total 2	p=3.7x10 <sup>-3</sup> *	p<0.001*	p<0.001*

I. Freshwater and marine; II: Brackish and freshwater species; III: Freshwater and terrestrial; IV: Freshwater and brackish water; V: Marine, freshwater and terrestrial; A. Dry season; B: Rainy season; FE: Fisher's exact test; FFH: Fisher-Freeman-Halton test; \*: significant difference ( $p<0.05$ )

Making a total of three species (7.5%) collected in the dry season and 17 species (42.5%) collected in the rainy season.




Based on the ecological impact, we recorded two species (5.0% of the species richness) potentially useful (*Ch. muelleri*)

(Chaetocerotaceae), and *Epithemia turgida* (Rhopalodiaceae)) ( $187.5 \times 10^5$  cells (1.7% of the global collection)). *Ch. muelleri* with  $62.5 \times 10^5$  cells (0.6%) is known as bio accumulator and detoxifier of antibiotics aquatic pollution, and *Ep. turgida* with  $125.0 \times 10^5$  cells (1.2%), through its N-fixing endosymbionts Cyanobacteria, can be used for the biofertilization of aquatic ecosystems. The two species were recorded exclusively at low tide in Kienke, the first in the dry season and the second in the rainy season. One species (2.5%) (*Ce. hirundinella* (Ceratidae)) known as harmful for fish, was rarely recorded at both tides ( $3.8 \times 10^5$  cells (0.9% at high tide and low tide respectively)) in the dry season and at low tide ( $25.0 \times 10^5$  cells (1.2%) in the rainy season in the Kienke). It is known to cause drastic blooms worldwide. The remaining 37 species (92.5%) were of unknown ecological impact apart from their photosynthetic ability placing them at the base of the fish food chain. These species were very abundant during both seasons (23 species (57.5%) in the dry season; 33 species (82.5%) in the rainy season; 37 species (92.5%) in the pooled seasons). It was the same in both river mouths (20 species (50.0%) in Nyong; 17 species (42.5%) in Kienke). Due to their preference for a specific habitat, they can be considered bio indicators of the water quality and food resource for fish. The weighed mean sensitivity of the taxa present in the collection (WWS) and the trophic diatom index (TDI) showed that in each river mouth, the community of diatoms were in the undisturbed conditions with no or little alteration of human origin and a low organic pollution (oligotrophic or mesotrophic state) (Nyong River mouth: WWS=3.1, TDI=52.7; Kienke River mouth: WWS=3.8, TDI=69.7; pooled rivers: WWS=3.6, TDI=65.0).

## 3.2. Alpha Diversity and the Community Structure

### 3.2.1. Alpha Diversity

The numbers of species recorded in both rivers at both tides and both seasons revealed a low species richness (richness ratio close to 0) (Table 3). The lowest species richness was noted in both rivers at high tide during the dry season (Nyong River mouth: 3 species; Margalef index:  $Mg=0.401$ ; richness ratio:  $d=0.020$ ; Kienke River mouth: 3 species,  $Mg=0.306$ ,  $d=0.004$ ).



**Figure 2.** Individual rarefaction curves of the Chromista collections during the dry season and the rainy season in Nyong and Kienke River mouths at both tides.

The highest species richness was recorded in the overall pooled data from both rivers and both seasons ( $S=40$  species;  $Mg=4.215$ ;  $d=0.004$ ) (Table 3). Other values of the species richness were found between the two extremes. Individual rarefaction curves approached the plateaus of saturation and for a standard sample of  $701.0 \times 10^5$  cells, the pooled data appeared the most species rich ( $E(S(n=701.0 \times 10^5))=39 \pm 1$  species), followed by the distribution in the rainy season ( $E(S(n=701.0 \times 10^5))=34 \pm 0$  species), and the assemblage in the dry season appeared less species rich ( $E(S(n=701.0 \times 10^5))=6 \pm 0$  species) (Figure 2A). A similar trend was noted in Nyong ( $E(S(n=131.0 \times 10^5))=19 \pm 1$  species for the pooled data;  $E(S(n=131.0 \times 10^5))=17 \pm 1$  species in the rainy season, and  $E(S(n=131.0 \times 10^5))=3 \pm 0$  species in the dry season) (Figure 2B), as well as in Kienke ( $E(S(n=671.0 \times 10^5))=20 \pm 0$  species for the pooled data;  $E(S(n=671.0 \times 10^5))=17 \pm 0$  species in the rainy season, and  $E(S(n=671.0 \times 10^5))=3 \pm 0$  species in the dry season) (Figure 2C).

In each river mouth, the species diversity recorded in rainy season was higher than that recorded in dry season (Table 3). Between the two river mouths, the species diversity recorded in Kienke was significantly higher than that recorded in the Nyong (Table 3). Both river mouths presented a high diversity (Shannon-Weaver's diversity values were close to the maximum value), a highly even community (Pielou's values were close to the unity) and a very low dominance by a few

species (Berger-Parker index were in all cases inferior to the median value and close to the null value; Table 3).

Pairwise comparison of diversity indexes showed significant differences (Table 3). Considering Chao1 index (non-parametric estimator of the 'TRUE' species richness), sampling success was maximal (100.0%) (Table 3), suggesting all rare species were collected. Although the rank-frequency plotting of the pooled SAD was close to the Fisher's log-series model (statistics:  $\alpha=5.261$ ,  $x=0.9995$ ,  $\chi^2=1.644 \times 10^4$ ,  $p=0.0$ ), a fairly weak concave appearance was noted, suggesting absence of a few co-dominants (Figure 3A). The similar shape was noted during the dry season ( $\alpha=0.8735$ ,  $x=0.999$ ,  $\chi^2=975.7$ ,  $p=3.4 \times 10^{-211}$ ; Figure 3B), and during the rainy season ( $\alpha=4.417$ ,  $x=0.9995$ ,  $\chi^2=1.473 \times 10^4$ ,  $p=0.0$ ; Figure 3C). A similar shape was noted during both seasons in Nyong (dry season:  $\alpha=0.534$ ,  $x=0.9964$ ,  $\chi^2=290.6$ ,  $p=3.7 \times 10^{-65}$ ; rainy season:  $\alpha=2.485$ ,  $x=0.9989$ ,  $\chi^2=4510.0$ ,  $p=0.0$ ; pooled seasons:  $\alpha=2.974$ ,  $x=0.9988$ ,  $\chi^2=5764.0$ ,  $p=0.0$ ; Figure 4), and during both seasons in Kienke (dry season:  $\alpha=0.4027$ ,  $x=0.9994$ ,  $\chi^2=1844.0$ ,  $p=0.0$ ; rainy season:  $\alpha=2.083$ ,  $x=0.9997$ ,  $\chi^2=1.118 \times 10^4$ ,  $p=0.0$ ; pooled seasons:  $\alpha=2.471$ ,  $x=0.9997$ ,  $\chi^2=1.315 \times 10^4$ ,  $p=0.0$ ; Figure 5).

Based on the Hill's  $N_1$  and  $N_2$  values, the numbers of abundant species were close to the number of co-dominants (Hill's ratio close to 1) (Table 3), suggesting a low dominance of the assemblages by a few species. On the base of the  $N_1$  diversity number (Table 3) and the rank-abundance plotting (Figures 3, 4 and 5), during the dry season, the recorded three species (*Cc. meneghiniana*, *De. elegans*, and *Op. cochleare*) were simply abundant and co-dominant species in the Nyong River mouth and three other species (*Ch. muelleri*, *Cr. erosa*, and *Su. linearis*) were simply abundant and co-dominants in Kienke (Table 3). During the rainy season, 14 species on the 17 recorded species (82.4%) (*Ac. exiguoides*, *Am. ovalis*, *An. sphaerophora*, *Au. granulate*, *Ca. noricus*, *Cm. solea*, *Co. placentula*, *Dp. arctica*, *Fa. construens*, *Fr. adnata*, *Gn. mutica*, *Ha. amphioxys*, *Pi. cardinaliculus*, and *Ta. flocculosa*) were simply abundant in the Nyong River mouth while 13 species on the 17 recorded species (76.5%) (*Cc. stelligera*, *Ce. hirundinella*, *De. thermalis*, *Dp. ovalis*, *Go. olivaceum*, *Gn. gigas*, *Cy. unicorn*, *Me. granulata*, *Ni. amphibia*, *Rh. longiseta*, *St. astraea*, *Su. capronii*, and *Sy. ulna*) were abundant in Kienke (Table 3).

**Table 3.** Matrix of the species richness, diversity, evenness and dominance indexes for the pooled data from both river mouths.

Indices	Nyong River mouths			Kienke River mouths			Both rivers		
	I	II	III	I	II	III	I	II	III
<b>A. Richness indexes</b>									
n x10 <sup>5</sup> cells	145.8	2,312.5	2,458.3	687.5	7,281.3	7,968.8	833.3	9,593.8	10,427.1
(%)	(1.4)	(22.2)	(23.6)	(6.6)	(69.8)	(76.4)	(8.0)	(92.0)	(100.0)
S (%)	3 (7.5)	17 (42.5)	20 (50.0)	3 (7.5)	17 (42.5)	20 (50.0)	6 (15.0)	34 (85.0)	40 (100.0)
n <sub>max</sub> x10 <sup>5</sup> cells	62.5	385.4	385.4	375.0	1,395.8	1,395.8	375.0	1,395.8	1,395.8
Magalef: Mg	0.401	2.065	2.433	0.306	1.799	2.115	0.743	3.599	4.215
Richness ratio: d = S/n	0.020	0.007	0.008	0.004	0.002	0.003	0.007	0.004	0.004
Chao1	3	17	20	3	17	20	6	34	40
% SE=(S/Chao 1)*100	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<b>B. Diversity indexes</b>									
Shannon-Weaver H'	1.004	2.612	2.742	0.918	2.530	2.685	1.398	3.102	3.245
H' <sub>max</sub> =ln(S)	1.099	2.833	2.996	1.099	2.833	2.996	1.792	3.526	3.689
Simpson's D	0.388	0.087	0.079	0.438	0.098	0.085	0.309	0.061	0.054
Hill's N <sub>1</sub> = e <sup>H'</sup>	2.730	13.629	15.522	2.503	12.553	14.654	4.048	22.250	25.656
Hill's N <sub>2</sub> = 1/D	2.579	11.447	12.719	2.286	10.213	11.766	3.236	16.276	18.525
Hill's ratio N <sub>2</sub> /N <sub>1</sub>	0.945	0.840	0.819	0.913	0.814	0.803	0.800	0.732	0.722
Rare species: Chao1-N <sub>1</sub>	0	3	4	0	4	5	2	12	14
<b>C. Evenness index</b>									

Indices	Nyong River mouths			Kienke River mouths			Both rivers		
	I	II	III	I	II	III	I	II	III
Pielou $J=(H'/H'_{\max})$	0.914	0.922	0.915	0.835	0.893	0.896	0.780	0.880	0.880
E. Dominance index									
$I_{BP} = n_{\max}/n$	0.429	0.166	0.156	0.545	0.192	0.175	0.449	0.145	0.134
Pairwise comparisons of diversity indexes (Student t-test)									
Comparison	Shannon-Weaver index $H'$					Simpson's diversity index			
I vs. II	Nyong: $t=-45.93$ ; $df=119.29$ ; $p=5.7 \times 10^{-108} *$ Kienke: $t=-76.82$ ; $df=961.07$ ; $p=0 *$					Nyong: $t=17.46$ ; $df=150.88$ ; $p=4.9 \times 10^{-38} *$ Kienke: $t=31.72$ ; $df=709.56$ ; $p=3.5 \times 10^{-138} *$			
Nyong vs. Kienke	Dry season: $t=-2.30$ ; $df=262.43$ ; $p=0.022 *$ Rainy season: $t=-5.28$ ; $df=4,285.20$ ; $p=1.3 \times 10^{-7} *$ Both seasons: $t=-3.62$ ; $df=4,316.60$ ; $p=3.0 \times 10^{-4} *$					Dry season: $t=2.47$ ; $df=273.73$ ; $p=0.014 *$ Rainy season: $t=4.46$ ; $df=4,615.20$ ; $p=8.3 \times 10^{-6} *$ Both seasons: $t=2.95$ ; $df=4,613.70$ ; $p=0.003 *$			

ns: not significant difference ( $p \geq 0.05$ ); \*: significant difference ( $p < 0.05$ ); n: sample size;  $n_{\max}$ : maximum abundance; S: observed species richness; Mg: Margalef richness index; d: richness ratio;  $H'$ : Shannon-Weaver diversity index;  $H_{\max}$ : maximum Shannon-Weaver diversity index; D: Simpson's diversity index;  $N_1$ : Hill's first order diversity number;  $N_2$ : Hill's second order diversity number; Hill: Hill's diversity ratio; J: Pielou's evenness index;  $I_{BP}$  = Berger-Parker dominance index; SE: sampling effort.

Making a total of 17 species (42.5%) in Nyong, 16 species (40.0%) in Kienke, and 33 species (82.5%) in the pooled distribution. Thirty-three species were represented by  $9,947.9 \times 10^5$  cells (95.4%) ( $145.8 \times 10^5$  cells (1.4%) in the dry season in Nyong,  $687.5 \times 10^5$  cells (6.6%) in the dry season in Kienke;  $2,166.7 \times 10^5$  cells (20.8%) in rainy season in Nyong;  $6,947.8 \times 10^5$  cells (66.8%) in rainy season in Kienke). The number rare species was low and they were scarce.

### 3.2.2. Adjustment of the SADs

Adjustment of SADs to five theoretical models showed a poor quality of fit (Pearson correlation  $r \leq -0.95$ ): dry season in both rivers:  $r=-0.884$ ,  $p=0.019$ , 6 species; dry season in Nyong:  $r=-0.867$ ,  $p=0.333$ , 3 species; dry season in Kienke:  $r=-0.993$ ,  $p=0.073$ , 3 species; rainy season in both rivers:  $r=-0.828$ ,  $p=1.5 \times 10^{-9}$ , 34 species; rainy season in Nyong:  $r=-0.912$ ,  $p=3.5 \times 10^{-7}$ , 17 species; rainy season in Kienke:  $r=-0.913$ ,  $p=3.3 \times 10^{-7}$ , 17 species; pooled seasons in both rivers:  $r=-0.817$ ,  $p=1.3 \times 10^{-10}$ , 40 species; pooled seasons in Nyong:  $r=-0.889$ ,  $p=1.6 \times 10^{-7}$ , 20 species; pooled seasons in Kienke:  $r=-0.896$ ,  $p=9.6 \times 10^{-8}$ , 20 species.


Based on AIC and BIC (Table 4) and SADs (Figures 3, 4 and 5), LN model fitted assemblages with high environmental constants ( $m' > 1$ ), except the pooled assemblage in dry season in both rivers which fitted ZM model. LN model fitted the pooled data with a low  $m'$  value (maximum abundance:  $n_1=1385.8 \times 10^5$  cells; sample size:  $n=10,427.1 \times 10^5$  cells; 40 species; deviance=151.06; mean of logarithms of abundance:  $x=2.206$ ; lognormal variance:  $\sigma^2=0.192$ ; lognormal standard deviation:  $\sigma=0.429$ ; environmental constant:  $m'=0.493$ ; correction factor: 1.049; corrected model:  $n_i=33,920.1 \times 10^5(0.350)^{P_i}$  with  $P_i$  as probits of species cumu-

lative percentage ranks (Table 4, Figure 3A).

This was the case of the pooled assemblage in the two rivers in the rainy season ( $n_1=1395.8 \times 10^5$  cells;  $n=9,593.8 \times 10^5$  cells; 34 species; deviance=107.92;  $x=2.252$ ;  $\sigma^2=0.18$ );  $\sigma=0.417$ ;  $m'=0.522$ ; correction factor: 1.050; corrected model:  $n_i=33,959.9 \times 10^5(0.358)^{P_i}$  (Table 4, Figure 3C). ZM model best fitted the pooled assemblage recorded during the dry season in the two river mouths with a low value of the average fractal dimension of the distribution of individuals among species ( $(1/\gamma) < 1$ ) (deviance=48.44,  $Q=833.3 \times 10^5$  cells,  $n_1=375.0 \times 10^5$  cells, 6 species; Gauss-Marquard's procedure for the determination of the model's parameters: starting point:  $x_0=(0; 2)^T$ ; tolerance of the functional value:  $\varepsilon=1.0 \times 10^{-10}$ ; damping factor:  $\lambda_0=100$ ; beta statistic:  $\beta=1.340$ ; gamma statistic:  $\gamma=1.093$ ; correction factor: 0.787; corrected model:  $n_i=656.0 \times 10^5(i+1.340)^{-1.093}$ ;  $1/\gamma=0.915$ ) (Table 4, Figure 3B). Assemblages in each season and each river fitted the LN model. In Nyong, it was the case of the assemblage in the dry season and the environmental constant was high ( $m' > 1.0$ ) ( $n_1=62.5 \times 10^5$  cells;  $n=145.8 \times 10^5$  cells; 3 species; deviance=9.46;  $x=1.637$ ;  $\sigma^2=0.076$ ;  $\sigma=0.248$ ;  $m'=1.469$ ; correction factor: 1.011; corrected model:  $n_i=2,862.8 \times 10^5(0.046)^{P_i}$  cells) (Table 4, Figure 4A), but the environmental constant was low in the rainy season ( $n_1=385.4 \times 10^5$  cells;  $n=2,312.5 \times 10^5$  cells; 17 species; deviance=154.94;  $x=2.029$ ;  $\sigma^2=0.103$ ;  $\sigma=0.309$ ;  $m'=0.947$ ; correction factor: 1.020; corrected model:  $n_i=6,296.1 \times 10^5(0.045)^{P_i}$  cells) (Table 4, Figure 4B). LN model fitted the pooled seasons with a low environmental constant ( $n_1=385.4 \times 10^5$  cells;  $n=2,458.3 \times 10^5$  cells; 20 species; deviance=53.20;  $x=1.970$ ;  $\sigma^2=0.115$ ;  $\sigma=0.326$ ;  $m'=0.853$ ; correction factor: 1.028; corrected model:



$n_i = 6,501.0 \times 10^5 (0.437)^{P_i}$  cells (Table 4, Figure 4C).



**Figure 3.** Rank-frequency diagram of the pooled Chromista collection in both river mouths at both tides, showing species in decreasing order of numerical occurrence.

In Kienke, assemblage in dry season fitted the LN model with a low environmental constant ( $n_i = 375.0 \times 10^5$  cells;  $n = 687.5 \times 10^5$  cells; 3 species; deviance=27.93;  $x = 2.256$ ;  $\sigma^2 = 0.167$ ;  $\sigma = 0.397$ ;  $m' = 0.574$ ; correction factor: 0.988; corrected model:  $n_i = 142,622.2 \times 10^5 (0.289)^{P_i}$  cells) (Table 4, Figure 5A). A similar result was obtained in rainy season ( $n_i = 1,395.8 \times 10^5$  cells;  $n = 7,281.3 \times 10^5$  cells; 17 species; deviance=94.08;  $x = 2.476$ ;  $\sigma^2 = 0.164$ ;  $\sigma = 0.400$ ;  $m' = 0.566$ ; correction factor: 1.001; corrected model:  $n_i = 56,730.5 \times 10^5 (0.358)^{P_i}$  cells) (Table 4, Figure 5B) as well as the pooled assemblage from both seasons ( $n_i = 1,395.8 \times 10^5$  cells;  $n = 7,968.8 \times 10^5$  cells; 20 species; deviance=109.54;  $x = 2.448$ ;  $\sigma^2 = 0.162$ ;  $\sigma = 0.395$ ;  $m' = 0.581$ ; correction factor: 1.008; corrected model:  $n_i = 46,270.4 \times 10^5 (0.367)^{P_i}$  cells) (Table 4, Figure 5C).

### 3.3. Beta Diversity: Dissimilarity Between SADs

Based on the species composition, a few common orders and families were recorded but no species was common to both rivers. In the pooled data, a null level of dissimilarity was detected between the two tides in the dry season (Table

5A) and it was the same in rainy season (Table 5A) as well as between tides in the pooled seasons (Table 5A). However, dissimilarity close to the median value was noted between the data set at high tide in the dry season (HS) and the pooled data at high tide in both rivers (Table 5A).

A low dissimilarity was noted between HS and the overall assemblage (Table 5A). Data at low tide in the dry season showed a null dissimilarity (Table 5A). In the rainy season, a null dissimilarity was noted between high tide and low tide (Table 5A). A low level of dissimilarity was recorded between high tide and the pooled assemblage at both tides in rainy season or the pooled assemblage at both tides and both seasons. A high level of dissimilarity was detected when compared to the assemblage at high tide in the pooled both seasons (Table 5A).

The assemblage at low tide showed a high dissimilarity to the assemblage at the pooled tides during the rainy season, to the assemblage at the low tide in the rainy season, or the pooled assemblage in the pooled seasons (Table 5A).

A low dissimilarity was noted between the pooled assemblage in the rainy season and the assemblage at high tide in the rainy season but it was high when compared to the as-


semblage at low tide in the pooled seasons or to the assemblage at the pooled tides in the pooled seasons (Table 5A). In the pooled data from both seasons, assemblage at high tide showed a null dissimilarity when compared to that noted at low tide. The dissimilarity was low when compared to the pooled tides (Table 5A). In the pooled seasons, low tide showed a high dissimilarity to the pooled tides (Table 5A). In Nyong, dissimilarity was null in both seasons between high and low tides (Table 5B), but in the dry season, it was high between the two tides and the pooled data (Table 5B). It was the same in the rainy season (Table 5B). Comparisons in the rainy season showed a null dissimilarity (Table 5B). Comparisons in the pooled seasons showed a low to null dissimilarity during the dry season and high dissimilarities in the rainy season (Table 5B).

Pooled data in rainy season and pooled seasons showed a high dissimilarity (Table 5B). In Kienke, the dissimilarity was null between the two tides in dry season (Table 5C) and

it was high when each tide was compared to the pooled data at both tides (Table 5C). Comparisons to data recorded during the rainy season showed a null dissimilarity (Table 5C). High tide in the pooled seasons showed a null dissimilarity when compared to low tide in each season, and the dissimilarity was low when compared to low tide in dry season or to the pooled tides in dry season (Table 5C). But the dissimilarity was high when compared to low tide in the rainy season or to the pooled tides in the same season (Table 5C).

### 3.4. Correlation Between Species

Twenty-one sample units (12 in Nyong, and nine in Kienke) allowed to detect an overall positive net association ( $VR > 1$ ) in Kienke (Schluter's Variance ratio:  $VR = 1.795$ ; statistic:  $W = 16.154$ ,  $df = 19$ ,  $p < 0.001$ ) and the pooled assemblage ( $VR = 1.043$ ;  $W = 21.91$ ,  $df = 20$ ,  $p < 0.001$ ).



**Figure 4.** Rank-frequency diagrams of the Chromista collection in the Nyong River mouth at both tides and both seasons, showing species in decreasing order of numerical occurrence.

In Nyong, a negative net association was noted ( $VR = 0.615$ ;  $W = 7.37$ ,  $df = 19$ ,  $p < 0.001$ ). Positive correlation

suggested that compared species were mutually tolerant (Table 6). In the pooled assemblage, correlations were positive (Table 6A). *Ac. exiguoides* was positively associated with *Co. placentula* (Table 6A). *Am. ovalis* was positively associated with *Cm. apiculata*, *Cs. rudolfi*, and *Gn. Mutica*. *Ca. noricus* was positively associated with the useful species *Ch. muelleri*. *Ce. hirundinella* was positively correlated with *Cr. erosa*, and *Cc. meneghiniana*. *Co. placentula* was positively correlated with *Cm. apiculata*, and *Gn. mutica* (Table 6A). *Cs. rudolfi* was positively correlated with *Fa. construens*. *Cr. erosa* was positively correlated with *Cc. meneghiniana*, *De. elegans*, and *Rh. longiseta* (Table 6A). *Cc. meneghiniana* was positively correlated with *Cc. stelligera* (Table 6A). *Cm. apiculata* was positively correlated with *Di. sertularia*, and *Gn. mutica* (Table 6A). *Cm. solea* was positively correlated with *De. elegans*, and *Ni. sigma* (Table 6A). *De. elegans* was positively correlated with one species (*Rh. longiseta*) (Table

6A). *De. thermalis* was positively correlated with *Gn. gigas*, and *Ni. tryblionella*. *Di. sertularia* was positively correlated with *Gn. mutica*, and *Ni. amphibia* (Table 6A). *Go. olivaceum* was positively correlated with *Rh. longiseta* (Table 6A). *Gn. mutica* was positively correlated with *Pi. cardinaliculus* (Table 6A). *Ni. amphibia* was positively correlated with *Ni. tryblionella* (Table 6A). *Pi. cardinaliculus* was positively correlated with *St. astraea* (Table 6A). In Nyong, correlations were positive (Table 6B). *Ac. exigoides*, *Cm. apiculata*, *Cm. solea*, *De. elegans* and *Gn. mutica* presented the correlation forms identical to those recorded in the pooled data (Table 6B).


In the case of *Am. ovalis*, the correlations in the pooled data existed in Nyong except that *Cs. rudolfi* in the pooled data was replaced by *Pi. cardinaliculus* in Nyong (Table 6B). *Ca. noricus* became positively correlated with *Cs. rudolfi* instead of *Ch. muelleri* as it was the case in the pooled data (Table 6B).

**Table 4.** Akaike Information Criteria (AIC) and Bayesian Information Criteria (BIC) values for the adjusted theoretical models.

AIC (BIC) and the best fitted model						
SAD model	Nyong River mouth			Kienke River mouth		
	Dry season 145.8x10 <sup>5</sup> cells 3 species	Rainy season 2,312.5x10 <sup>5</sup> cells 17 species	Pooled data 2,457.3x10 <sup>5</sup> cells 20 species	Dry season 687.5x10 <sup>5</sup> cells 3 species	Rainy season 7,281.3x10 <sup>5</sup> cells 17 species	Pooled data 7,968.8x10 <sup>5</sup> cells 20 species
Broken-Stick (BS)	37.34 (37.34)	299.05 (299.05)	307.81 (307.81)	45.25 (45.25)	245.70 (245.70)	319.54 (319.54)
Log-linear (LL)	34.99 (34.06)	178.43 (179.26)	220.08 (221.08)	58.29 (57.39)	283.22 (284.06)	369.49 (370.48)
Log-normal (LN)	30.32 (28.51) *	153.27 (154.94) *	184.82 (186.81) *	53.04 (51.24) *	226.26 (227.92) *	262.82 (264.81) *
Zipf (Z)	34.21 (32.41)	214.60 (216.27)	252.86 (254.85)	77.00 (75.20)	627.03 (628.70)	688.24 (690.23)
Zipf-Mandelbrot (ZM)	32.32 (29.61)	153.67 (156.17)	186.35 (189.34)	55.04 (52.34)	266.80 (269.30)	347.75 (350.74)

AIC (BIC) and the best fitted model			
SAD model	Pooled rivers		
	Dry season n=833.3x10 <sup>5</sup> cells S=6 species	Rainy season n=9,593.8x10 <sup>5</sup> cells S=34 species	Pooled data n=10,427.1x10 <sup>5</sup> cells S=40 species
Broken-Stick (BS)	106.49 (106.49)	529.39 (529.39)	630.97 (630.97)
Log-linear (LL)	93.65 (93.44)	583.01 (584.53)	718.18 (719.87)
Log-normal (LN)	96.72 (96.30)	350.85 (353.90) *	431.95 (435.33) *
Zipf (Z)	109.78 (109.37)	845.13 (848.18)	1,039.65 (1043.03)
Zipf-Mandelbrot (ZM)	92.41 (91.79) *	579.59 (584.17)	NA

AIC: Akaike information criteria; BIC: Bayesian information criteria; \* best fitted theoretical model in bold; NA: Not available



**Figure 5.** Rank-frequency diagrams of the Chromista collection in the Kienke River mouth at both tides and both seasons, showing species in decreasing order of numerical occurrence.

**Table 5.** Bray-Curtis dissimilarity values between data sets recorded at two tides during two seasons in the Nyong and Kienke river mouths.

Dry season			Rainy season			Pooled seasons		
High tide	Low tide	Pooled tides	High tide	Low tide	Pooled tides	High tide	Low tide	Pooled tides
A. Overall data sets								
Dry season								
High tide	1.000							
Low tide	0.0	1.000						
Pooled tides	1.000	0.0	1.000					
Rainy season								
High tide	0.0	0.0	0.0	1.000				
Low tide	0.0	0.0	0.0	0.0	1.000			
Pooled tides	0.0	0.0	0.0	0.313	0.898	1.000		
Pooled seasons								
High tide	0.431	0.0	0.431	0.841	0.0	0.295	1.000	
Low tide	0.0	0.0	0.0	0.0	1.000	0.898	0.0	1.000
Pooled tides	0.123	0.0	0.123	0.295	0.864	0.966	0.386	0.864
B. Nyong River mouth								



	Dry season			Rainy season			Pooled seasons		
	High tide	Low tide	Pooled tides	High tide	Low tide	Pooled tides	High tide	Low tide	Pooled tides
Dry season									
High tide	1.000								
Low tide	0.0	1.000							
Pooled tides	0.727	0.600	1.000						
Rainy season									
High tide	0.0	0.0	0.0	1.000					
Low tide	0.0	0.0	0.0	0.0	1.000				
Pooled tides	0.0	0.0	0.0	0.600	0.727	1.000			
Pooled seasons									
High tide	0.145	0.0	0.137	0.959	0.0	0.585	1.000		
Low tide	0.0	0.087	0.082	0.0	0.977	0.715	0.0	1.000	
Pooled tides	0.066	0.050	0.113	0.574	0.699	0.969	0.608	0.721	1.000
C. Kienke River mouth									
Dry season									
High tide	1.000								
Low tide	0.0	1.000							
Pooled tides	0.706	0.625	1.000						
Rainy season									
High tide	0.0	0.0	0.0	1.000					
Low tide	0.0	0.0	0.0	0.0	1.000				
Pooled tides	0.0	0.0	0.0	0.151	0.957	1.000			
Pooled seasons									
High tide	0.558	0.0	0.453	0.760	0.0	0.144	1.000		
Low tide	0.0	0.086	0.081	0.0	0.977	0.936	0.0	1.000	
Pooled tides	0.090	0.076	0.159	0.139	0.913	0.955	0.217	0.935	1.000

**Table 6.** Kendall's tau correlation recorded between *Chromista* species in 21 sample units from the Nyong and Kienke River mouths.

Species 1/Species 2	$\tau$ (p-value)	Species 1/Species 2	$\tau$ (p-value)	Species 1/Species 2	$\tau$ (p-value)
A. Overall pooled data from both seasons and both river mouths (n=21)					
<i>Ac. exiguoides</i>		<i>Cs. rudolfi</i>		<i>De. thermalis</i> (continued)	
<i>Co. placentula</i>	0.737(3.0x10-6) *	<i>Fa. construens</i>	0.447(0.005) *	<i>Ni. tryblionella</i>	0.425(0.007) *
<i>Am. ovalis</i>		<i>Cr. erosa</i>		<i>Di. sertularia</i>	
<i>Cm. apiculata</i>	0.474 (0.003) *	<i>Cc. meneghiniana</i>	0.445 (0.005) *	<i>Gn. mutica</i>	0.364 (0.021) *
<i>Cs. rudolfi</i>	0.322 (0.041) *	<i>De. elegans</i>	0.445 (0.005) *	<i>Ni. amphibia</i>	0.568 (3.2x10-4) *
<i>Gn. mutica</i>	0.742(2.5x10-6) *	<i>Rh. longiseta</i>	0.360 (0.023) *	<i>Go. olivaceum</i>	
<i>Ca. noricus</i>		<i>Cc. meneghiniana</i>		<i>Rh. longiseta</i>	0.536 (0.001) *

Species 1/Species 2	$\tau$ (p-value)	Species 1/Species 2	$\tau$ (p-value)	Species 1/Species 2	$\tau$ (p-value)
<i>Ch. muelleri</i>	0.520 (0.001) *	<i>Cc. stelligera</i>	0.497 (0.002) *	<i>Gn. mutica</i>	
<i>Ce. hirundinella</i>		<i>Cm. apiculata</i>		<i>Pi. cardinaliculus</i>	0.645 (4.4x10 <sup>-5</sup> ) *
<i>Cr. erosa</i>	0.431 (0.006) *	<i>Di. sertularia</i>	0.568(3.2x10 <sup>-4</sup> )*	<i>Ni. amphibia</i>	
<i>Cc. meneghiniana</i>	0.378 (0.016) *	<i>Gn. mutica</i>	0.716(5.6x10 <sup>-6</sup> )*	<i>Ni. tryblionella</i>	0.533 (0.001) *
<i>Ch. muelleri</i>		<i>Cm. solea</i>		<i>Pi. cardinaliculus</i>	
<i>Go. olivaceum</i>	0.424 (0.007) *	<i>De. elegans</i>	0.645(4.4x10 <sup>-5</sup> )*	<i>St. astraea</i>	0.424 (0.007) *
<i>Ni. amphibia</i>	0.592(1.7x10 <sup>-4</sup> ) *	<i>Ni. sigma</i>	0.716(5.6x10 <sup>-6</sup> )*		
<i>Rh. longiseta</i>	0.332 (0.035) *	<i>De. elegans</i>			
<i>Co. placentula</i>		<i>Rh. longiseta</i>	0.379 (0.016) *		
<i>Cm. apiculata</i>	0.598(1.5x10 <sup>-4</sup> ) *	<i>De. thermalis</i>			
<i>Gn. mutica</i>	0.385 (0.015) *	<i>Gn. gigas</i>	0.328 (0.037) *		
B. Nyong River mouth (n=12)					
<i>Ac. exiguoides</i>		<i>Cs. rudolfi</i> (continued)		<i>Cm. solea</i>	
<i>Co. placentula</i>	0.647 (0.006) *	<i>Fa. construens</i>	0.671 (0.004) *	<i>De. elegans</i>	0.580 (0.013) *
<i>Am. ovalis</i>		<i>Cr. erosa</i>		<i>Ni. sigma</i>	0.725 (0.002) *
<i>Cm. apiculata</i>	0.580 (0.013) *	<i>De. elegans</i>	0.725 (0.002) *	<i>De. elegans</i>	
<i>Gn. mutica</i>	0.895 (1.3x10 <sup>-4</sup> ) *	<i>Ni. amphibia</i>	0.580 (0.013) *	<i>Rh. longiseta</i>	0.725 (0.002) *
<i>Pi. cardinaliculus</i>	0.725 (0.002) *	<i>Rh. longiseta</i>	0.474 (0.043) *	<i>Di. sertularia</i>	
<i>Ca. noricus</i>		<i>Cc. stelligera</i>		<i>Gn. mutica</i>	0.474 (0.043) *
<i>Cs. rudolfi</i>	0.580 (0.013) *	<i>Cm. solea</i>	0.725 (0.002) *	<i>Fa. construens</i>	
<i>Ce. hirundinella</i>		<i>Cm. apiculata</i>		<i>Ni. amphibia</i>	0.671 (0.004) *
<i>De. elegans</i>	0.671 (0.004) *	<i>Di. sertularia</i>	0.725 (0.002) *	<i>Gn. mutica</i>	
<i>Co. placentula</i>		<i>Gn. mutica</i>	0.725 (0.002) *	<i>Pi. cardinaliculus</i>	0.580 (0.013) *
<i>Cm. apiculata</i>	0.620 (0.008) *	<i>Cm. apiculata</i>			
<i>Cs. rudolfi</i>		<i>Rh. longiseta</i>	0.580 (0.013) *		
<i>Di. sertularia</i>	0.580 (0.013) *				
C. Kienke River mouth (n=9)					
<i>Am. ovalis</i>		<i>Cs. rudolfi</i>		<i>Di. sertularia</i>	
<i>Cs. rudolfi</i>	1.000 (1.7x10 <sup>-4</sup> ) *	<i>De. thermalis</i>	0.730 (0.006) *	<i>Ni. amphibia</i>	0.548 (0.040) *
<i>De. thermalis</i>	0.730 (0.006) *	<i>Cr. erosa</i>		<i>Ni. tryblionella</i>	1.000 (1.7x10 <sup>-4</sup> ) *
<i>Ca. noricus</i>		<i>Rh. longiseta</i>	0.553 (0.038) *	<i>Go. olivaceum</i>	
<i>Ch. muelleri</i>	1.000 (1.7x10 <sup>-4</sup> ) *	<i>Cc. stelligera</i>		<i>Rh. longiseta</i>	0.789 (0.003) *
<i>Ni. amphibia</i>	0.730 (0.006) *	<i>Di. sertularia</i>	0.730 (0.006) *	<i>Ni. amphibia</i>	
<i>Ce. hirundinella</i>		<i>Ni. tryblionella</i>	0.730 (0.006) *	<i>Pi. cardinaliculus</i>	0.548 (0.040) *
<i>St. astraea</i>	0.617 (0.021) *	<i>De. thermalis</i>			
<i>Ch. muelleri</i>		<i>Di. sertularia</i>	0.548 (0.040) *		
<i>Ni. amphibia</i>	0.730 (0.006) *	<i>Ni. tryblionella</i>	0.548 (0.040) *		

*Ce. hirundinella* was positively correlated with *De. elegans* (Table 6B). *Co. placentula* was correlated with *Cm. apiculata* (Table 6B). *Cs. rudolfi* was positively correlated with *Di. sertularia* in Nyong (Table 6B). In the case of *Cr. erosa*, correlations noted in the pooled data existed in Nyong except *Cc. meneghiniana* in the pooled data which was replaced by *Ni. amphibia* in Nyong (Table 6B). The correlations in the pooled data concerning *Cc. meneghiniana*, *Cm. solea*, *De. thermalis*, *Go. olivaceum*, *Ni. amphibia*, and *Pi. cardinaliculus* were no longer found in Nyong (Table 6B). *Di. sertularia* was positively correlated with *Gn. mutica* (Table 6B).

In Kienke, correlations were positive (Table 6C). *Ca. noricus* was positively correlated with *Ni. amphibia* (Table 6C). *Ce. hirundinella* was positively correlated with *St. astraea* (Table 6C). *Ch. muelleri* was positively correlated with *Ni. amphibia* (Table 6C). *Cs. rudolfi* was positively correlated with *De. thermalis* (Table 6C). *Cr. erosa* was positively correlated with *Rh. longiseta* (Table 6C). *Cc. stelligera* and *De. thermalis* were each positively correlated with *Di. sertularia* and *Ni. tryblionella* (Table 6C). *Di. sertularia* was positively correlated with *Ni. amphibia* and *Ni. tryblionella*. *Go. olivaceum* was positively correlated with *Rh. longiseta* and *Ni. amphibia* was positively correlated with *Pi. cardinaliculus* (Table 6C). Other correlations were not significant.

## 4. Discussion

### 4.1. Species Richness and Diversity of Chromista

The present study is the first step in an in-depth study of the algae assemblage in Nyong and Kienke (Southern Cameroon), evaluating the place occupied by zoonotic species, or those useful for the nutrition of fish. Collected cells belonged to three phyla, eight classes, 23 orders, 32 genera and 40 species [20] species (50.0%) from each river mouth) and no common species was recorded. Melosiraceae was the most collected family (13.4%), followed by Bacillariaceae (12.2%), Surirellaceae (10.8%), Gomphonemataceae (9.6%), Stephanodiscaceae (7.7%), Goniochloridaceae (7.2%), Rhizosoleniaceae (7.0%), Fragilariaceae (6.3%), Cryptomonadaceae (3.5%), Ceratiaceae (3.0%), Catenulaceae (2.8%), Phytodiniaceae (2.6%), Aulacoseiraceae (2.2%), Diploneidaceae (1.9%), Cocconeidaceae (1.4%), Achnanthaceae (1.2%), Anomoeoneidaceae (1.2%), Rhopalodiaceae (1.2%) while other families were rare and represented each by less than 1.0% of the total collection. The most specious family was Bacillariaceae (15.0% of the total species richness), followed by Surirellaceae (12.5%), Stephanodiscaceae (7.5%). Diploneidaceae and Fragilariaceae were each represented by two species (5.0%). Other families were rare. The high presence and abundance of diatoms is not surprising, as it is the case worldwide, in stagnant or slow flowing waters [43, 48].

Based on the information on the recorded contributing diatom species [64-66], the weighed mean sensitivity of taxa (WWS) and the trophic diatom index (TDI) showed that in the studied rivers, the diatom community was in the reference conditions (undisturbed) with no or little alteration of human origin and a low organic pollution (oligotrophic or in mesotrophic state) [13-22]. Values were relatively close to 56.74 reported in Lake Porsuk Dam in Turkey the value [65] and 39 to 44.5 reported in the mesotrophic waters of Upper Porsuk Creek Kütahya in Turkey [67]. *Me. granulata* was the most collected (13.4%), followed by *Go. olivaceum* (9.6%), *Rh. longiseta* (7.0%), *Gn. gigas* (6.4%), *De. thermalis* and *Sy. ulna* (5.1% respectively), *St. astraea* (3.9%), *Ca. noricus* (3.7%), 3.6% respectively for *Cr. erosa* and *Ni. amphibia*, *Cc. stelligera* (3.2%), *Ce. hirundinella* (3.0%), *Am. ovalis* (2.8%), *Cy. unicorn* (2.6%), *Su. linearis* (2.4%), *Su. capronii* (2.3%), *Au. granulata* (2.2%), *Ha. amphioxys* (2.1%). Other species were rare. Twenty-four freshwater species [60.0%: 12 species (30.0%) in Nyong and Kienke respectively] were recorded. Three marine species (7.5%: one species *Cs. rudolfi* in Nyong and two species *Ch. muelleri*, and *Ep. turgida* in Kienke) were recorded. One brackish species (2.5%) (*De. thermalis*) was noted exclusively in Kienke. Twelve tolerant species (30.0%) (seven (17.5%) in Nyong and five (12.5%) in Kienke) were recorded. Two potentially useful species (the antibiotics pollution detoxifier *Ch. muelleri* [21, 24] and the biofertilizer *Epithemia turgida* [23]) were recorded. One species harmful to the fish was recorded (the bloom forming *Ce. hirundinella* known as fish-killer species [31]) and 37 species (92.5%) of unknown toxigenic status were able to be considered as bioindicators. These waters allowed the development of autotrophic and/or mixotrophic algae species. Useful species were few, lowly abundant and masked by indifferent species which were good resource of the nutrition for aquatic macroinvertebrates and/or macrovertebrates. The recorded harmful species is known to form blooms in stagnant waters of lakes, ponds and reservoirs as well as in slow-moving freshwaters, when water temperatures are warmer than usual [31] as the case in Cyanobacteria [31, 41, 51, 66, 67]. In the studied mouths, the harmful species found came in addition to the presence of 16 species of toxigenic Cyanobacteria [41], reinforcing the poor quality of the raw waters for human drinking and for fish farming. Non-toxicogenic Chromista are abundant and diverse, suggesting either the continuous re-colonization from the tributary rivers, or an appearance of tolerant species adapted to the unstable conditions. Adaptation is well illustrated in tolerant species [11, 27, 52]. Twelve tolerant species (30.0%) were recorded (seven (17.5%) in Nyong and five (12.5%) in Kienke) and the two rivers presented a low species richness compared to the situation in other freshwaters. For example, 11 classes, 45 genera and 75 species were reported in Mezam River (Bamenda, Cameroon), amongst which Bacillariophyceae was the most represented class and Naviculaceae was the dominant family [68]. A total of 237 epilithic diatom

taxa, mostly cosmopolitan, belonging to 39 genera distributed among 25 families were reported in Mfoundi River (Yaounde, Cameroon) [69]. It is possible that patterns in Nyong and Kienke depended on local environmental conditions or the sampling methodology and design. Useful species and harmful species were lowly recorded probably because of the regulation of their populations by competing species and/or local natural enemies. Their density could increase if the environmental balance is disrupted due to the climate change force and the growing anthropization actions.

## 4.2. Community Structure and Functioning Model

Both rivers presented high species diversity, high even community and a low dominance by a few species. In Nyong, the dissimilarity was null in both seasons at high and low tides. In Kienke, the dissimilarity was null at the two tides in dry season and was high when each tide was compared to the pooled tides. The overall assemblage showed a negative global net association in Nyong and a positive net association in Kienke and in the pooled distribution. In each river, assemblage fitted the LN model with low environmental constants ( $m' < 1$ ) except in the dry season in Nyong where  $m'$  was high. Pooled SAD fitted in the dry season, the ZM model with a low fractal dimension of the distribution of individuals among species ( $(1/\gamma) < 1$ ). LN model describes a community where the majority of species show moderate abundances and it is reported fitting SADs of zooplankton in coastal neritic and estuarine conditions in the Arcachon Bay (France) [70], the freshwater snails at swampy areas in Douala (Cameroon) [71], the Cyanobacteria in Nyong and Kienke [41]. Given that nomocenosis are associations of species subject to the influence of the same factors, they characterize evolved or less disturbed environments. ZM model is frequently adapted to evolved communities in natural environments, with a multi-species network structure corresponding to an optimal structure for the circulation of information [55, 72, 73]. The pooled assemblage functioned in dry season on the basis of maintaining the complex information network developed at spatio-temporal scales (close to the ecological balance) with a low significant force of regeneration compared to undisturbed freshwaters.

## 5. Conclusion

The aim of the study was to establish a baseline of information on the distribution of chromists, in order to evaluate the status and the occurrence of species known as bio-indicators of the aquatic life quality. The studied river mouths presented two useful species and one harmful species to fish. A total of  $10,427.1 \times 10^5$  cells belonged to three phyla, eight classes, 23 orders, 32 genera and 40 species. Bacillariophyta was the most collected phylum (71.7%). Melosiraceae was the most collected family (13.4%). *Melosira granulata*

was the most collected species (13.4%). The species diversity was high. Assemblages were highly even and lowly dominated by a few species. Twenty-four freshwater species (60.0%) were recorded. Three marine species (2.4%) were recorded and one brackish specialist was recorded exclusively in Kienke (5.1%). Thirteen tolerant species (32.6%) were recorded. The trophic diatom index showed that the diatoms were in undisturbed conditions. The useful species and the harmful species were rare and exclusively found in Kienke. Species exhibited a positive net association in Kienke and the pooled assemblage from both rivers in the rainy season while it was negative in Nyong. Assemblages fitted LN model with low environmental constants ( $m' < 1$ ) except in dry season in Nyong where  $m'$  was high suggesting that majority of species were moderately abundant (evolved environments). Pooled assemblage in the dry season functioned on the basis of maintaining a complex information network developed at spatio-temporal scales with a low force of regeneration.

## Abbreviations

- AIC: Akaike Information Criteria  
*Ac. exiguoides*: *Achnanthes exiguoides* Comp ère, 1967  
*Am. ovalis*: *Amphora ovalis* (K ütz ing) K ütz ing, 1844  
*An. sphaerophora*: *Anomoeoneis sphaerophora* E. Pfitzer, 1871  
*Au. granulata*: *Aulacoseira granulata* (Ehrenberg) Simonsen, 1979  
BIC: Bayesian Information Criteria  
BS: Broken-Stick Theoretical Model (McArthur's model)  
*Ca. noricus*: *Campylodiscus noricus* Ehrenberg ex K ütz ing, 1844  
*Cc. meneghiniana*: *Cyclotella meneghiniana* f. *meneghiniana* K ütz ing, 1844  
*Cc. stelligera*: *Cyclotella stelligera* Cleve & Grunow, 1882  
*Ch. muelleri*: *Chaetoceros muelleri* Lemmermann, 1898  
*Ce. hirundinella*: *Ceratium hirundinella* (O. F. Müller) Dujardin, 1841  
*Cm. apiculata*: *Cymatopleura solar* var. *apiculata* W. Smith, 1853  
*Cm. solea*: *Cymatopleura solea* var. *baicalensis* Skvortzow & Meyer, 1928  
*Co. placentula*: *Cocconeis placentula* Ehrenberg, 1838  
*Cr. erosa*: *Cryptomonas erosa* Ehrenberg 1832  
*Cs. rudolfi*: *Coscinodiscus rudolfi* Bachmann, 1938  
*Cy. unicorn*: *Cystodinium unicorn* G. A. Klebs, 1912  
*De. elegans*: *Denticula elegans* var. *elegans* K ütz ing, 1844  
*De. thermalis*: *Denticula thermalis* var. *fossilis* Frenguelli, 1936  
*Di. sertularia*: *Dinobryon sertularia* Ehrenberg, 1834  
*Dp. arctica*: *Diploneis arctica* (Lange–Bertalot) Lange–Bertalot & A. Fuhrmann, 2016  
*Dp. ovalis*: *Diploneis ovalis* var. *pumila* (Grunow) Cleve, 1894



*Ep. turgida*: *Epithemia turgida* var. *turgida* (Ehrenberg) Kützinger, 1844

*Fa. construens*: *Fragilaria construens* f. *construens* (Ehrenberg) Grunow, 1862

*Fr. adnata*: *Frustulia adnata* Kützinger, 1833

*Gn. gigas*: *Goniochloris gigas* Bourrelly

*Gn. mutica*: *Goniochloris mutica* (A. Braun) Fott, 1960

*Go. olivaceum*: *Gomphonema olivaceum* (Lyngbye) Desmazières, 1825

*Ha. amphioxys*: *Hantzschia amphioxys* (Ehrenberg) Grunow, 1880

LL: Lognear

LN: Lognormal

*Ma. smithii*: *Mastogloia smithii* Thwaites ex W. Smith, 1856

*Me. granulata*: *Melosira granulata* var. *angustissima* Otto Müller, 1899

*Ni. amphibia*: *Nitzschia amphibia* Grunow, 1862

*Ni. sigma*: *Nitzschia sigma* (Kützinger) W. Smith, 1853

*Ni. tryblionella*: *Nitzschia tryblionella* Hantzsch, 1860

*Op. cochleare*: *Ophiocyrtium cochleare* (Eichwald) A. Braun 1855

*Pi. cardinaliculus*: *Pinnularia cardinaliculus* var. *ceylonica* Skvortzow, 1930

*Rh. longiseta*: *Rhizosolenia longiseta* O. Zacharias, 1893

*St. astraea*: *Stephanodiscus astraea* (Ehrenberg) Grunow, 1880

*Su. capronii*: *Surirella capronii* Brébisson & Kitton, 1869

*Su. linearis*: *Surirella linearis* f. *kolhapurensis* Sarode & Kamat, 1984

*Sy. ulna*: *Synedra ulna* (Nitzsch) Ehrenberg, 1832

SAD(s): Species Abundance Distribution(s)

sp.: species plurimae

*Ta. flocculosa*: *Tabellaria flocculosa* (Roth) Kützinger, 1844

VR: Variance Ratio of Schluter

Z: Zipft

ZM: Zipft-Mandelbrot

## Acknowledgments

The authors acknowledge Mrs. Fumdjum Marie for financing laboratory analyzes and the staff the micro-algae laboratory of the Specialized Center for Research on Marine Ecosystems at Kribi (AquaSol service) for the hospitality. They thank canoeists for making their canoe available, elders of Laboratory of Biotechnology and Environment (University of Yaounde 1) and the Laboratory of Biology of Animal Organisms (University of Douala) for assistance in species identification, data analysis and the manuscript preparation.

## Author Contributions

**Christelle Chimene Mokam:** Conceptualization, Data

curation, Formal Analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing

**Andrea Sarah Kenne Toukem:** Software, Formal Analysis, Writing – original draft

**Christian Dongmo Teufack:** Investigation, Methodology

**Fabien Tresor Amougou Dzou:** Investigation, Methodology

**Sedrick Junior Tsekane:** Software, Formal Analysis, Writing – original draft

**Mohammadou Moukhtar:** Formal Analysis, Writing – original draft

**Auguste Pharaon Mbianda:** Software, Formal Analysis, Writing – original draft

**Martin Kenne:** Conceptualization, Resources, Data curation, Software, Formal Analysis, Supervision, Funding acquisition, Validation, Writing – original draft, Project administration, Writing – review & editing

## Funding

The Cameroonian Ministry of Higher Education provided funds through the research support program.

## Data Availability Statement

The data supporting the outcome of this research work has been reported in this manuscript. They are available from the corresponding author upon reasonable request.

## Conflicts of Interest

The authors declare no conflicts of interest.

## References

- [1] Guiry, M. D. How many species of algae are there? A reprise. Four kingdoms, 14 phyla, 63 classes and still growing. *Journal of Phycology*. 2024, 00, 1–15. <https://doi.org/10.1111/jpy.13431>
- [2] Cavalier-Smith, T. Eukaryote kingdoms: seven or nine? *Bio-systems*. 1981, 14, 461-481. [https://doi.org/10.1016/0303-2647\(81\)90050-2](https://doi.org/10.1016/0303-2647(81)90050-2)
- [3] Cavalier-Smith, T. Early evolution of eukaryote feeding modes, cell structural diversity, and classification of the protozoan phyla Loukozoa, Sulcozoa, and Choanozoa. *European Journal of Protistology*. 2013, 49, 115-178. <https://doi.org/10.1016/j.ejop.2012.06.001>
- [4] Cavalier-Smith, T. Kingdom Chromista and its eight phyla: a new synthesis emphasising periplastid protein targeting, cytoskeletal and periplastid evolution, and ancient divergences. *Protoplasma*. 2018, 255, 297-357. <https://doi.org/10.1007/s00709-017-1147-3>
- [5] Aubert, D. Une nouvelle mégaclassification pragmatique du vivant. *Médecine/Sciences*. 2016, 32, 497-499. <https://doi.org/10.1051/medsci/20163205017>

- [6] Johnson, L. R., John, D. M., Whitton, B. A., Brook, A. J. Phylum Xanthophyta (Tribophyta) (Yellow-Green Algae). In *The Freshwater Algal Flora of the British Isles: An Identification Guide to Freshwater and Terrestrial Algae*, Second Edition, Johnson, L. R., Ed., Cambridge University Press. 2021, pp. 318-345.  
<https://doi.org/10.1017/CHOL9781108784122.019>
- [7] Guiry, M. D. How many species of algae are there? *Journal of Phycology*. 2012, 48(5), 1057–1063.  
<https://doi.org/10.1111/j.1529-8817.2012.01222.x>
- [8] Ruggiero, M. A., Gordon, D. P., Orrell, T. M., Bailly, N., Bourgoin, T., Brusca, R. C., Cavalier-Smith, T., Guiry, M. D., and, Kirk, P. M. A Higher Level Classification of All Living Organisms. *PLoS ONE*. 2015, 10(4), e0119248.  
<https://doi.org/10.1371/journal.pone.0119248>
- [9] Hagen, J. Five Kingdoms, More or Less: Robert Whittaker and the Broad Classification of Organisms. *BioScience*, 2012, 62. 67-74. <https://doi.org/10.1525/bio.2012.62.1.11>
- [10] Integrated Taxonomic Information System. “Chromista”. Available from [www.itis.gov](http://www.itis.gov), CC0). [Accessed 29 February 2024], <https://doi.org/10.5066/F7KH0KBK>
- [11] DiatomBase. “Chromista”. Available from <https://diatombase.org> [Accessed 26 February 2024].
- [12] Bloch, K., Vardi, P. Microalgae as photosynthetic oxygen generators for pollution control, life support systems and medicine, In *Algae: Nutrition, Pollution Control and Energy Sources*, Hagen, K. N., Ed., Nova Science Publishers, Inc. Tel-Aviv University, Petah Tikva, Israel; 2009, pp. 3-11.
- [13] Zalat, A. A., Nitychoruk, J., Chodyka, M., Pidek, I. A., Welc, F. Recent and fossil freshwater diatoms of Poland: taxonomy, distribution and their significance in the environmental reconstruction. Part 1. Coscinodiscophyceae, Mediophyceae and Fragilariophycidae. John Paul II University of Applied Sciences, Biała Podlaska; 2022, pp. 1-306
- [14] Jahn, R., Kusber, W. H., Romero, O. E. *Cocconeis pediculus* Ehrenberg and *C. placentula* Ehrenberg var. *placentula* (Bacillariophyta): Typification and taxonomy. *Fottea*. 2009, 9(2), 275–288. <https://doi.org/10.5507/fot.2009.027>
- [15] Hecky, R. E., Kilhan, P. Diatoms in Alkaline, Saline Lakes: Ecology and Geochemical Implications. *Limnology and Oceanography*. 1973, 18(1), 53-71.  
<https://doi.org/10.4319/LO.1973.18.1.0053>
- [16] Spaulding, S. A., Potapova, M. G., Bishop, I. W., Lee, S. S., Gasperak, T. S., Jovanovska, E., Edlund, M. B.. Diatoms.org: supporting taxonomists, connecting communities. *Diatom Research*. 2021, 36(4): 291-304.  
<https://doi.org/10.1080/0269249X.2021.2006790>
- [17] Houk, V.; Klee, R., Tanaka, H. Atlas of freshwater centric diatoms with a brief key and descriptions, Part III, Stephanodiscaceae A, Cyclotella, Tertarius, Discostella. *Fottea*. 2010, 10(Supplement), 1–498.
- [18] Krammer, K., Lange-Bertalot, H. Bacillariophyceae, 2nd part: Bacillariaceae, Epithemiaceae, Surirellaceae. In *Süßwasserflora von Mitteleuropa*, Band 2/2. Ettl, H., Gerloff, J., Heynig, H., Mollenhauer, D., Eds., Gustav Fischer Verlag, Stuttgart, Germany; 1988, pp. 1-596.
- [19] Krammer, K., Lange-Bertalot, H. Bacillariophyceae, 3. Teil: Centrales, Fragilariaceae, Eunotiaceae. In *Süßwasserflora von Mitteleuropa*, Band 2/3. Ettl, H., Gerloff, J., Heynig, H., Mollenhauer, D., Eds., Gustav Fischer Verlag, Stuttgart, Germany; 1991, pp. 1-596.
- [20] Mohanty, T. R., Tiwari, N. K., Kumari, S., Ray, A., Manna, R. K., Bayen, S., Roy, S., Gupta, S. D., Ramteke, M. H., Swain, H. S., Bhor, M., Das, B. K. Variation of *Aulacoseira granulata* as an eco-pollution indicator in subtropical large river Ganga in India: a multivariate analytical approach. *Environmental science and pollution research international*. 2022, 29(25), 37498-37512  
<https://doi.org/10.1007/s11356-021-18096-9>
- [21] Taylor, J. C., Harding, W. R., Archibald, C. G. M. An Illustrated Guide to Some Common Diatom Species from South Africa. WRC Report TT 282/07. Water Research Commission. Gezina, Pretoria, South Africa; 2007, pp. 1-225.
- [22] DeColibus, D. “*Tabellaria flocculosa* (Roth) Kütz. 1844. In *Diatoms of North America*”. Available from <https://diatoms.org> [Accessed 01 March 2024].
- [23] Lowe, R. “*Epithemia turgida*. In *Diatoms of North America*”. Available from [https://diatoms.org/species/epithemia\\_turgida](https://diatoms.org/species/epithemia_turgida) [Accessed 22 January 2024].
- [24] Mojiri, A., Baharlooeian, M., Zahed M. A. The Potential of *Chaetoceros muelleri* in Bioremediation of Antibiotics: Performance and Optimization. *International Journal of Environmental Research in Public Health*. 2021, 22, 18(3), 977.  
<https://doi.org/10.3390/ijerph18030977>
- [25] Sivonen, K. Cyanobacterial toxins and toxin production. *Phycologia*. 1996, 35, 12-24.  
<https://doi.org/10.2216/I0031-8884-35-6S-12.1>
- [26] Mankiewicz, J., Tarczyska, M., Walter, Z., Zalewski, M. Natural Toxins from Cyanobacteria, *Acta Biologica Cracoviensia Series Botanica*. 2003, 45(2), 9-20.
- [27] WHO (World Health Organization). Chapter 8 – Algae and Cyanobacteria in Freshwater. In *Guidelines for safe recreational waters. Volume 1 – Coastal and fresh waters*. Geneva, Switzerland; 2003, pp. 136-158.
- [28] Zanchett, G., Oliveira-Filho, E. Cyanobacteria and Cyanotoxins: From Impacts on Aquatic Ecosystems and Human Health to Anticarcinogenic Effects. *Toxins*. 2013, 5, 1896-917.  
<https://doi.org/10.3390/toxins5101896>
- [29] Rastogi, R. P., Madamwar, D., Incharoensakdi, A. Bloom Dynamics of Cyanobacteria and Their Toxins: Environmental Health Impacts and Mitigation Strategies. *Frontiers in Microbiology*. 2015, 6, 1254. <https://doi.org/10.3389/fmicb.2015.01254>
- [30] Beasley, V. R. Harmful Algal Blooms (Phycotoxins). In *Reference Module in Earth Systems and Environmental Sciences*, Elsevier Inc. 2019.  
<https://doi.org/10.1016/B978-0-12-409548-9.11275-8>

- [31] Nicholls, K. H., Kennedy, W., Hammett, C. A fish-kill in Heart Lake, Ontario, associated with the collapse of a massive population of *Ceratium hirundinella* (Dinophyceae). *Freshwater Biology*. 1980, 10, 553-561.  
<https://doi.org/10.1111/j.1365-2427.1980.tb01231.x>
- [32] Caturao, R. D. Harmful and toxic algae. In *Health Management in Aquaculture* (2<sup>nd</sup> edition). Aquaculture Department, Southeast Asian Fisheries Development Center, Lio-Po G. D. & Inui, Y., Eds., Tigbauan, Iloilo, Philippines; 2010, pp. 170-182.
- [33] Magray, A. R., Lone, S. A., Ganai, B. A., Ahmad, F., Hafeez, S. The first detection and in vivo pathogenicity characterization of *Saprolegnia delica* from Kashmir Himalayas. *Aquaculture*. 2021, 542, 736876.  
<https://doi.org/10.1016/j.aquaculture.2021.736876>
- [34] Raghukumar, S. Ecology of the marine protists, the Labyrinthulomycetes (Thraustochytrids and Labyrinthulids). *European Journal of Protistology*. 2022, 38(2), 127-145.  
<https://doi.org/10.1078/0932-4739-00832>
- [35] Brummett, R. E., Nguenga, D., Tiotsop, F., Abina, J.-C... The commercial fishery of the middle Nyong River, Cameroon: productivity and environmental threats. *Smithiana Bulletin*. 2010, 11, 3-16.
- [36] Olivry, J. C. Monographie hydrologique. Fleuves et rivières du Cameroun. Ministère de l'Enseignement Supérieur et de la Recherche Scientifique au Cameroun - Institut Français de Recherche Scientifique pour le Développement en Coopération/ (MESRES-ORSTOM). Collection "Monographies Hydrologiques ORSTOM" No 9, Paris. 1986, pp. 1-781.
- [37] Dounias, E., Cogels, S., Mvé Mbida, S., Carrière, S. The safety net role of inland fishing in the subsistence strategy of multiactive forest dwellers in southern Cameroon. *Revue d'Ethnoécologie*. 2016, 10, 11-56.
- [38] Mama, A. C., Mbeng, O., Dongmo, C., Ohondja, A. Tidal Variations and its impacts on the abundance and diversity of phytoplankton in the Nyong estuary of Cameroon. *Journal of Multidisciplinary Engineering Science and Technology*. 2016, 3(1), 3159-3199.
- [39] Mama, A. C., Youbouni Ghepdeu, G. F., Ngoupayou Ndam, J. R., Bonga, M. D., Onana, F. M., Onguene, R. Assessment of water quality in the lower Nyong estuary (Cameroon, Atlantic Coast) from environmental variables and phytoplankton communities' composition. *African Journal of Environmental Science and Technology*. 2018, 12(6), 198-208.  
<https://doi.org/10.5897/AJEST2017.2454>
- [40] Essomba Biloa, R. E., Noah Ewoti, O. V., Tuekam Kayo, R. P., Sob Nangou, P. B., Tchakounté S., Onana, F. M., Nyamsi Tchatcho, N. L., Zebaze Togouet S. H. Zooplankton Dynamics of the Kienke Estuary (Kribi, South Region of Cameroon): Importance of Physico-Chemical Parameters. *Open Journal of Ecology*. 2021, 11, 837-869.  
<https://doi.org/10.4236/oje.2021.1112051>
- [41] Mokam, C. C., Kenne Toukem, A. S., Dongmo Teufack, C., Amougou Dzou, F. T., Tsekane, S. J., Moukhtar, M., Mbianda, A. P., Kenne M. Water Quality, Biodiversity and Abundance of Blue-Green Algae in Nyong and Kienke River Mouths (South-Cameroon). *International Journal of Ecotoxicology and Ecobiology*. 2024, 9(1), 1-27.  
<https://doi.org/10.11648/j.ijee.20240901.11>
- [42] Climate-Data.org. Climat Kribi (Cameroun). Available from <https://fr.climate-data.org> [Accessed 22 January 2024].
- [43] Compère, P. Algues de la région du Lac Tchad. IV: Diatomophycées. *Cahier ORSTOM, série. Hydrobiologie*. 1975, 9(4), 205-290.
- [44] Carty, S. *Freshwater Dinoflagellates of North America*. Comstock Publishing Associates, Cornell University Press. Ithaca, New York, USA. 2014, pp. 280.  
<https://doi.org/10.7591/9780801470370>
- [45] Olenina, I., Hajdu, S., Edler, L., Andersson, A., Wasmund, N., Busch, S., Göbel, J., Gromisz, S., Huseby, S., Huttunen, M., Jaanus, A., Kokkonen, P., Ledaine, I., Niemkiewicz, E.. Biovolumes and size-classes of phytoplankton in the Baltic Sea HELCOM Balt. Sea Environ. Proc. No. 106. 2006, pp. 1-144.
- [46] Bellinger, E. G., Sigee, D. C. *Freshwater Algae. Identification and Use as Bioindicators*. Wiley-Blackwell, Hoboken, USA. 2010, pp. 1-271.
- [47] Jahn, R., Kusber, W.-H. A Key to Diatom Nomenclature. *Diatom Research*. 2009, 24(1), 101-111.  
<https://dx.doi.org/10.1080/0269249X.2009.9705785>
- [48] DREAL (Direction Régionale de l'Environnement, de l'Aménagement et du logement d'Occitanie). "Atlas des Diatomées de l'ex partie Languedoc-Roussillon. Languedoc Roussillon". Available from <https://www.occitanie.developpement-durable.gouv.fr/atlas-des-diatomees-de-l-ex-partie-languedoc-a25336.html> [Accessed 22 January 2021].
- [49] Sonneman, J. A., Sincok, A., Fluin, J., Reid, M., Newall, P., Tibby, J., Gell, P. An Illustrated Guide to Common Stream Diatom Species from Temperate Australia. Cooperative Research Centre for Freshwater Ecology. Identification guide n°33. Land & Water Resources Research & Development Corporation. 2<sup>nd</sup> Australian Algal Workshop, Adelaide University, 17-19th April, 2000, pp. 1-166.
- [50] Kumar, D. S., Maurya, O. N. Floristic Survey of Algae in Vikramsila Gangetic Dolphin Sanctuary, Bihar (India). *Nelumbo*. 2015, 57, 124-134.  
<https://doi.org/10.20324/nelumbo/v57/2015/87116>
- [51] WoRMS (World Register of Marine Species). "Chromista". Available from <https://www.marinespecies.org/aphia.php?p=taxdetails&id=7> at VLIZ. [Accessed 06 February 2024].
- [52] Guiry, M. D., Guiry, G. M. 2024. "Kingdom *Chromista* Cavalier-Smith 1981". *AlgaeBase*. World-wide electronic publication, University of Galway, Ireland. Available from <https://www.algaebase.org> [Accessed 28 February 2024].

- [53] Wilson, J. B. Methods for fitting dominance/diversity curves. *Journal of Vegetation Science*. 1991, 2(1), 35-46. <https://doi.org/10.2307/3235896>
- [54] Iganaki, H. Mise au point de la loi de Motomura et essai d'une ecologie évolutive. *Vie et Milieu*. 1967, 18, 153-166.
- [55] Li, W. Zipf's Law Everywhere. *Glottometrics*. 2002, 5, 14-21.
- [56] Zipf, G. K.,. Human Behaviour and the Principle of Least Effort: An introduction to human ecology. 2nd edition, Hafner, New York, NY, USA. 1965, pp. 1-573.
- [57] Le, D.-H., Pham, C.-K., Nguyen, T. T. T., Bui, T. T. Parameter extraction and optimization using Levenberg-Marquardt algorithm. In *Proceedings of 2012 IEEE conference. Fourth International Conference on Communications and Electronics (ICCE)*, Hanoi University of Science and Technology, Hanoi (Vietnam), 2022; pp. 434-437. <https://doi.org/10.1109/CCE.2012.6315945>
- [58] Frontier, S. Applications of Fractal Theory to Ecology. In *Developments in Numerical Ecology.*, Legendre P., Legendre, L., Eds., NATO ASI Series book series (volume 14). Springer, Berlin, Heidelberg. 1987, pp. 335-378.
- [59] Bach, P., Amanieu, M., Lam-Hoai, T., Lasserre, G. Application du modele de distribution d'abondance de Mandelbrot a l'estimation des captures dans l'étang de Thau. *Journal du Conseil/Conseil Permanent International pour l'Exploration de la Mer*. 1988, 44, 235-246. <https://doi.org/10.1093/icesjms/44.3.235>
- [60] Murthy, Z. V. P. Nonlinear Regression: Levenberg-Marquardt Method, In *Encyclopedia of Membranes*, Drioli, E., Giorno L., Eds., Springer-Verlag, Berlin, Heidelberg, 2014, pp. 1-3.
- [61] Ludwig, J. A., Reynolds, J. F. *Statistical Ecology*. John Wiley & Sons, New York, USA. 1988, pp. 1-337.
- [62] Friedman, N. & Potapova, M. "*Pinnularia cardinaliculus*". In *Diatoms of North America*. Available from <https://diatoms.org> [Accessed 22 January 2024].
- [63] Lange-Bertalot, H., Fuhrmann, A. Contribution to the genus *Diploneis* (Bacillariophyta): Twelve species from Holarctic freshwater habitats proposed as new to science. *Fottea*. 2016, 16(2), 157-183. <https://doi.org/10.5507/fot.2015.027>
- [64] Kelly, M. G., Whitton, B. A.. The trophic diatom index: a new index for monitoring eutrophication in rivers. *Journal of Applied Phycology*. 1995, 7, 433-444. <https://doi.org/10.1007/BF00003802>
- [65] Tokatli, C. Evaluation of water quality by using trophic diatom index: example of Porsuk Dam Lake. *Journal of Applied Biological Sciences* 2013, 7(1), 01-04.
- [66] Kelly, M. G., Adams, C., Graves, A. C., Jamieson, J., Krokowski, J., Lycett, E. B., Murray-Bligh, J., Pritchard, S., Wilkins, C. The trophic diatom index: A User's manual. Revised Edition. R & D technical Report E2/TR2. Environment Agency, Almondsbury, Bristol, BS32 4UD; 2001, pp. 1-135.
- [67] Solak, C., N. The Application of Diatom Indices in the Upper Porsuk Creek Kütahya – Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*. 2011, 11, 31-36. <https://doi.org/10.4194/tjfas.2011.0105>
- [68] Ndjouondo, G. P., Nwamo, R. D., Choula Tegantchouang, F. Diversity and Structure of Microalgae in the Mezam River (Bamenda, Cameroon). *African Journal of Environment and Natural Science Research*. 2023, 6(1), 19-35. <https://doi.org/10.52589/AJENS-R09H3LUP0>
- [69] Menye, D. É., Zébazé Togouet, S. H., Menbohan, S. F., Kemka, N., Nola, M., Boutin, C., Nguetso, V. F., Djaouda, M., Njiné T. Bio-écologie des diatomées épilithiques de la rivière Mfoundi (Yaoundé Cameroun): diversité distribution spatiale et influence des pollutions organiques. *Revue des sciences de l'eau/Journal of Water Science*. 2012, 25(3), 203-218. <https://doi.org/10.7202/1013103ar>
- [70] Castel, J., Courties, C. Composition and differential distribution of zooplankton in Arcachon Bay. *Journal of Plankton Research*. 1982, 4(3), 417-433. <https://doi.org/10.1093/plankt/4.3.417>
- [71] Kenne, E. L., Yetchom-Fondjo, J. A., Moumite, M. B., Tsekane, S. J., Ngamaleu-Siewe, B., Fouelifack-Nintidem, B., Biawa-Kagmegni, M., Tuekam Kowa, P. S., Fantio, R. M., Yomon, A. K., Kentsop-Tsafong, R. M., Dim-Mbianda, A. M., Kenne M. Species composition and diversity of freshwater snails and land snails at the swampy areas and streams edges in the urban zone of Douala, Cameroon. *International Journal of Biological and Chemical Sciences*. 2021, 15(4), 1297-1324. <https://dx.doi.org/10.4314/ijbcs.v15i4.2>
- [72] Shea, K., Chesson, P. Community ecology theory as a framework for biological invasions. *Trends in Ecology and Evolution*. 2002, 17(4), 170-176. [https://doi.org/10.1016/S0169-5347\(02\)02495-3](https://doi.org/10.1016/S0169-5347(02)02495-3)
- [73] Ferreira, F. C., Petrere-Jr., M. Comments about some species abundance patterns: classic, neutral, and niche partitioning models, *Brazilian Journal of Biology*, 2008, 68(4, Suppl.), 1003-1012. <https://doi.org/10.1590/S1519-69842008000500008>